



Received: 01-07-2024  
Accepted: 11-08-2024

## International Journal of Advanced Multidisciplinary Research and Studies

ISSN: 2583-049X

### New Asian "Jumping Worm" Records (Oligochaeta: Megascolecidae) in Ontario, Canada

John Warren Reynolds

Oligochaetology Laboratory, 9-1250 Weber Street East, Kitchener, ON Canada N2A 4E1, and Research Associate, New Brunswick Museum, Saint John, NB Canada

DOI: <https://doi.org/10.62225/2583049X.2024.4.4.3148>

Corresponding Author: **John Warren Reynolds**

#### Abstract

This is the first record of megascolecid "jumping worms" from Wellington County, Ontario. *Amyntas agrestis* has previously been reported from Essex, Wentworth and York Counties, and *Amyntas tokioensis* from Chatham-Kent and

Wentworth Counties in Ontario. Finding megascolecids under sod in the middle of a lawn is believed to be a second report.

**Keywords:** Oligochaeta, Megascolecidae, *Amyntas agrestis*, *Amyntas tokioensis*, under sod

#### Introduction

The Asian "jumping worms" (Oligochaeta: Megascolecidae) are expanding their range in Canada. Although there are 16 species<sup>H</sup> reported in North America north of Mexico, only five species have been reported to date in Canada (Chang *et al.*, 2016)<sup>[3]</sup>. The first literature report for Canada was from the Ojibway Prairie in Essex County, Ontario (*Amyntas agrestis* and *Metaphire hilgendorfi* as *Amyntas hilgendorfi*) (Reynolds, 2014)<sup>[26]</sup>. The second was also from Ontario: Chatham-Kent Co., Wheatley (*Amyntas tokioensis*, *Metaphire hilgendorfi*), Wentworth Co., Dundas (*Amyntas agrestis*, *Amyntas tokioensis*), and York Co. (East York, Toronto) (*Amyntas agrestis*, *Metaphire hilgendorfi*, *Pithemera bicincta*) (Reynolds and McTavish, 2021)<sup>[29]</sup>. The following year pheretimoid earthworms were collected in New Brunswick, York County (Burton) (*Amyntas minimus*, *Amyntas tokioensis*, *Metaphire hilgendorfi*) (McAlpine *et al.*, 2022)<sup>[16]</sup>. Earlier this year, *Amyntas tokioensis* was reported from Saint-Jerôme, Region of Montréal, Québec (Moore and Reynolds, 2024)<sup>[19]</sup>.

I have been studying them in Canada for over a decade, although my first research on these species was in the southeastern United States in the late 1960s (Reynolds, 1978)<sup>[23]</sup>. At that time, very few collections of megascolecids from natural habitats were found north of Virginia (Reynolds, 2010)<sup>[24]</sup>. There were a few reports from New England, in greenhouses and golf courses (Gates, 1954, 1958; Grant, 1955a,b,c)<sup>[5, 6, 9, 10, 11]</sup>.

There have been citizen reports for years, claiming they are present in various regions. During some of my lectures this spring, several participants told me they had "jumping worms" in areas where pheretimoids had not been previously recorded on their property and/or in their fields. I told them I would come and collect samples when they informed me that adults were present. Since these species overwinter as cocoons, it was necessary to give the cocoons time to mature for reliable identification. Fortunately, for these property owners the earthworms in their fields and gardens turned out to be the normal European Lumbricidae found throughout Canada (Reynolds, 2024)<sup>[28]</sup>.

<sup>H</sup> *Amyntas agrestis* (Goto and Hatai, 1899)<sup>[8]</sup>, *A. carnosus* (Goto and Hatai, 1899)<sup>[8]</sup>, *A. corticis* (Kinberg, 1867)<sup>[14]</sup>, *A. gracilis* (Kinberg, 1867)<sup>[14]</sup>, *A. hupeiensis* (Michaelsen, 1895)<sup>[18]</sup>, *A. loveridgei* (Gates, 1968)<sup>[7]</sup>, *A. minimus* (Horst, 1893)<sup>[13]</sup>, *A. morrisi* (Beddard, 1892)<sup>[1]</sup>, *A. rodericensis* (Grube, 1879)<sup>[12]</sup>, *A. tokioensis* (Beddard, 1892)<sup>[1]</sup>, *Metaphire californica* (Kinberg, 1867)<sup>[14]</sup>, *M. hilgendorfi* (Michaelsen, 1892)<sup>[17]</sup>, *M. houlleti* (Perrier, 1872)<sup>[21]</sup>, *M. posthuma* (Vaillant, 1868)<sup>[31]</sup>, *Pithemera bicincta* (Perrier, 1875)<sup>[22]</sup>, *Polypheretima elongata* (Perrier, 1872)<sup>[21]</sup>. Those in **bold** have been previously reported in Canada.


Citizen reports on sites like *iNaturalist* may or may not be accurate. Frequently, the photograph does not show an external character needed for identification to species. In the case of the Asian "jumping worms" (Megascolecidae) and the North American native species of *Diplocardia* (Acanthodrilidae) internal dissection is required to confirm the species. To remedy this, some of us have set up a program to help improve the situation (Mathieu *et al.*, 2023) [15].

### Objectives of the Study

- X Examine the range of pheretimoid earthworms in Ontario
- X Identify the specimens collected to species level
- X Record detailed habitat data.

### Methods

The earthworms were dug with a pitch fork from compost piles, under debris and sod where the property owners have collected "jumping worms" this year. Collection data included location, date, co-ordinates, soil temperature, elevation and collectors (Fig 1). The samples were brought back to the Oligochaetology Laboratory, preserved and identified to species.



**Fig 1:** The locations (M) and 3 of the habitat photos in Puslinch Township, Wellington County, Ontario, Canada. A B *Amyntas* in garden bed, B B *Amyntas* in compost pile, C B *Amyntas* under sod.

### Results

#### Material examined:

Ontario, Wellington Co., Puslinch Township, 2744 Concession 11, 43. 32.196E, -80 00.000E; elevation 2241 metres asl, soil temperature at 5 cm 24EC, flower garden, under sod, etc., canopy over the grassy areas white pine (*Pinus strobus* L.) and birch (*Betula papyrifera* Marshall) with maples (*Acer* spp.) in the fence rows, 3 August 2024, collectors George and Diane Tyszka with John W. Reynolds.

*Amyntas agrestis* 0-0-34 (Fig 2), *Amyntas tokioensis* 0-0-9 (Fig 3).



**Fig 2:** Dorsal view of *Amyntas agrestis*



**Fig 3:** Lateral view of *Amyntas tokioensis*

For age classification formula explanation, see Reynolds *et al.* (1974) [30] or Reynolds (2022) [27].


### Discussion

These specimens represent the first record of Asian "jumping worms" from Wellington County, Ontario. The two species, *Amyntas agrestis* and *A. tokioensis* have been reported previously from Ontario including the adjacent county of Wentworth (Reynolds and McTavish, 2021) [29]. Recently, there have been unconfirmed reports that pheretimoid worms may be present in the Arboretum at the University of Guelph (Wellington County). Over a decade ago, when I collected at that location, no pheretimoid ("jumping worms") were found (Reynolds, 2011) [25].

I have been collecting these megascolecid worms for more than 55 years and this is the first time I have seen or heard of them being under sod in a residential grass lawn. From my own collections and other literature reports, *Amyntas agrestis* and other megascolecid species have been found in gardens where commercial mulch has been used (Reynolds, 2014; Bellitürk *et al.*, 2015) [26, 2]. This is usually the method of spreading these earthworms (Nouri-Aiin *et al.*, 2022) [20]. My first collections of the megascolecids were in the Cherokee National Forest in east Tennessee (Reynolds, 1978) [23]. Subsequently, *A. agrestis* and others have been reported from deciduous forests of the mid-Atlantic USA (Chang *et al.*, 2017) [4]. *Amyntas agrestis*, *A. tokioensis* and *Metaphire hilgendorfi* are frequently found together in the same habitat (Chang *et al.*, 2016) [3]. Two or three of these species have co-habited in other Canadian provinces, Ontario (Reynolds and McTavish, 2021) [29] and New Brunswick (McAlpine *et al.*, 2022) [16]. The recent first

record of a pheretimoid earthworm in Quebec yielded only *A. tokioensis* (Moore and Reynolds, 2024) [18].

The current distribution of pheretimoid earthworms ("jumping worms") is shown in Fig 4 (below). There are numerous reports of these worms in other parts of Ontario but they are only referred to as "jumping worm(s)" without genus or species designation. Hopefully, in the future samples will be obtained by an oligochaetologist, so that we can produce a complete range of these worms in Ontario and Canada.



**Fig 4:** Location of pheretimoid collections in Ontario, Canada. A B Ojibway Prairie Provincial Nature Reserve (Essex Co.), B B Wheatley (Chatham-Kent Co.), C B Dundas (Hamilton, Wentworth Co.), D B East York (Toronto, York Region), E B Toronto, F B Blythwood Ravine (Toronto), S B Fonthill (Niagara Region) and Toronto (York Region), S B suspected species, unconfirmed. Species collection sites: M *Amyntas agrestis*, M *Amyntas hilgendorfi*, M *Amyntas tokioensis*, M *Pithemera bicincta*. New records G B Puslinch (Wellington Co.) (modified from Reynolds and McTavish, 2021) [29].

### Conflict of Interest

The author declares no conflict of interest.

### Funding

This research received no external funding.

### Conclusions

This study extends the range of pheretimoid earthworms in Ontario, including the first record of two species in Wellington County, *Amyntas agrestis* and *A. tokioensis*. This is believed to be the second report of these earthworms living under lawn sod as opposed to their usual habitat in forests, flower/ornamental gardens or cultivated fields.

### Acknowledgements

I wish to thank Wilma M. Reynolds of the Oligochaetology Laboratory (Kitchener, Ontario) for reviewing the manuscript, her comments and suggestions. Also, my appreciation extends to Bruce Miller of Elmira, Ontario for graphic assistance.

### References

1. Beddard FE. On some species of the genus *Perichaeta* (sensu stricto). Proceedings of the Zoological Society

London 1892: 153B172. [*Amyntas morrisoni* n. sp., *A. tokioensis* n. sp.], 1892.

2. Bellitürk K, Görres JH, Kunkle J, Melnichuk RDS. Can commercial mulches be reservoirs of invasive earthworms? Promotion of ligninolytic enzyme activity and survival of *Amyntas agrestis* (Goto and Hatai, 1899). Applied Soil Ecology. 2015; 87:27B31.
3. Chang C-H, Snyder BA, Szilávecz K. Asian pheretimoid earthworms in North America north of Mexico: An illustrated key to the genera *Amyntas*, *Metaphire*, *Pithemera*, and *Polypheretima* (Clitellata: Megascolecidae). Zootaxa. 2016; 4179(3):495B529.
4. Chang C-H, Szilávecz K, Buyer JS. *Amyntas agrestis* invasion increases microbial biomass in Mid-Atlantic deciduous forests. Soil Biology and Biochemistry. 2017; 14:189B199.
5. Gates GE. Exotic earthworms in the United States. Bulletin of the Museum of Comparative Zoology, Harvard. 1954; 111:219B258.
6. Gates GE. On some species of the oriental earthworm genus *Pheretima* Kinberg, 1867, with a key to species reported from the Americas. American Museum Novitates, No. 1888: 1B33, 1958.
7. Gates GE. On an anthropochorous species of the earthworm genus *Pheretima* (Megascolecidae, Oligochaeta). Journal of the Natural History 2: 253B261. [*Amyntas loveridgei* n. sp.], 1968.
8. Goto S, Hatai S. New or imperfectly known species of earthworms. No. 2. Annotationes Zoologicae Japonenses. 1899; 3(1):13B24. [*Amyntas agrestis* n. sp.]
9. Grant WC, Jr. Studies on moisture relationships in earthworms. Ecology. 1955a; 36(3):400B407.
10. Grant WC, Jr. Temperature relationship in the megascolecid earthworm *Pheretima hupeiensis*. Ecology. 1955b; 36(3):412B417.
11. Grant WC, Jr. An anatomical study of the peregrine megascolecid earthworm *Pheretima hupeiensis* in the eastern United States. Proceedings of the United States National Museum. 1955c; 105(3351):49B63.
12. Grube AE. Annelida. Philosophical Transactions of the Royal Society of London. 1879; 168:554B556. [*Amyntas rodericensis* n. sp.]
13. Horst R. Earthworms from the Malay Archipelago. Zoologische Ergebnisse einer reise in Niederländisch Ost-Indien. 1893; 3:28B77. [*Amyntas minimus* n. sp.]
14. Kinberg JG. Annulata nova. [transl. New Annelida.] Öfversigt af Kongl. Vetenskaps-akademiens förhandlingar Stockholm. 1867; 23:97B103, 356B357. [*Amyntas aeruginosus* n. gen. and n. sp., *Amyntas corticis* n. sp., *A. gracilis* n. sp.; *Metaphire californica* n. sp.]
15. Mathieu J, Bartz M, Brown G, Cameron E, Decaens T, Dupont L, Reynolds JW *et al.* Collaboration call: Building an image collection for computer vision identification of worldwide earthworm species in iNaturalist, 2023. <https://www.inaturalist.org/projects/curated-collection-of-earthworms-id>
16. McAlpine DF, Reynolds JW, Manzer L, Elton K. First reports of invasive earthworms (Oligochaeta: Megascolecidae) of Asian origin in Atlantic Canada. BioInvasions Records. 2022; 11(4):830B838.

17. Michaelsen W. Terricolen der Berliner Zoologischen Sammlung. II. [transl. Terricoles of the Berlin Zoological Collection. II.] Archiv für Naturgeschichte. 1892; 58(2):209B261. [*Amyntas hilgendorfi* n. sp.]
18. Michaelsen W. Zur Kenntnis der Oligochaeten. [transl. On the knowledge of oligochaetes.] Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg. 1895; 13(2):37. [*Amyntas hupeiensis*, n. sp.]
19. Moore J-D, Reynolds JW. First mention of exotic Asian earthworms in the Quebec Province, Canada. BioInvasions Records. 2024; 13:1B8.
20. Nouri-Aiin M, Connolly S, Keough C, Smigelsky AJ, Wen Y, Howland J, et al. Genetic population structure and reproductive system of two invasive Asian earthworms, *Amyntas tokioensis* and *Amyntas agrestis*. PeerJ. 2022; 10:e13622. Doi: <http://doi.org/10.7717/peerj.13622>
21. Perrier E. Recherches pour servir à l'histoire des Lombriciens terrestres. [transl. Research to serve the history of terrestrial earthworms.] Nouvelles Archives du Muséum d'Histoire Naturelle, Paris. 1872; 8:5B198. [*Metaphire houlleti* n. sp.; *Polypheretima elongata* sp. n.]
22. Perrier E. Sur les vers de terre des îles Philippines et de la Cochinchine. [transl. On the earthworms of the Philippine Islands and Cochinchina.] Comptes rendus hebdomadaires des Séances de l'Académie des Sciences, Paris (D). 1875; 81:1043B1046. [*Pithemera bicincta* n. sp.]
23. Reynolds JW. The earthworms of Tennessee Oligochaeta). IV. Megascolecidae, with notes on distribution, biology and a key to the species in the state. Megadrilogica. 1978; 3(7):117B129.
24. Reynolds JW. The earthworms (Oligochaeta: Acanthodrilidae, Lumbricidae, Megascolecidae and Sparganophilidae) of northeastern United States, revisited. Megadrilogica. 2010; 14(7):101B157.
25. Reynolds JW. Earthworms of the University of Guelph Arboretum. Megadrilogica. 2011; 15(2):15B20.
26. Reynolds JW. A checklist by counties of earthworms (Oligochaeta: Lumbricidae, Megascolecidae and Sparganophilidae) in Ontario, Canada. Megadrilogica. 2014; 16(10):111B135.
27. Reynolds JW. The Earthworms (Lumbricidae, Megascolecidae and Sparganophilidae) in Canada. Ottawa: Canada Food Inspection Agency, 2022, i + 179.
28. Reynolds JW. New lumbricid records (Oligochaeta: Lumbricidae) in Ontario, Canada while searching for the invasive "jumping worms" (Megascolecidae). GSC Advanced Research and Reviews. 2024; 20(2):16B20.
29. Reynolds JW, McTavish MJ. New Asian pheretimoid "jumping earthworm" records (Oligochaeta: Megascolecidae) in Canada. Megadrilogica. 2021; 26(7):83B89.
30. Reynolds JW, Clebsch EEC, Reynolds WM. The earthworms of Tennessee (Oligochaeta). I. Lumbricidae. Contribution to North American Earthworms, No. 3, Bulletin of Tall Timbers Research Station, No. 17, 1974; viii + 133 pp.
31. Vailliant L. Note sur l'anatomie de deux espèces du genre *Perichaeta* et essai de classification des Annélides Lombricines. [transl. Note on the anatomy of two species of the genus *Perichaeta* and attempt at classification of the Lombricine Annelids.] Mémoires de l'Académie des Sciences et Lettres de Montpellier. 1868; 7:143B173. [*Metaphire posthuma* n.sp.]

**Corresponding author:**

Dr. John Warren Reynolds

john.w.reynolds1941@gmail.com

<https://orcid.org/0000-0001-8362-9071>