



Received: 07-12-2025
Accepted: 17-12-2025

International Journal of Advanced Multidisciplinary Research and Studies

ISSN: 2583-049X

Letter to the Editor

Carotid Intima-Media Thickness and Periventricular White Matter Hyperintensities are Unlikely Predictors of Cognitive Decline

Josef Finsterer

Department of Neurology, Neurology & Neurophysiology Center, Vienna, Austria

Corresponding Author: **Josef Finsterer**

Letter to the Editor

We read with interest the article by Li *et al.* on a prospective cohort study on the association between carotid intima-media thickness (CIMT) assessed by ultrasound and periventricular white matter hyperintensities (PWMH) assessed by 3T NRI with mild cognitive impairment (MCI) assessed by 10 neuropsychological tests across 5 domains in 541 patients with Parkinson's disease ^[1]. The prevalence of MCI was 45% and independent predictors of MCI were PWMHs, CIMT, homocysteine levels, Hoehn and Yahr stage, and MDS-UPDRS score ^[1]. The diagnostic value was higher for combined CIMT and PWMHs compared to either marker alone ^[1]. The study is appealing but several points merit discussion.

The first point is that CIMT varies greatly between examinations and between observers ^[2]. The level at which CIMT was measured in the common carotid artery, the bulb, and the internal carotid artery was not defined ^[1]. Since the thickening of the artery wall varies greatly depending on the measurement height, it is important to measure CIMT at the same height in all included patients. The authors should also indicate the inter- and intra-observer variability of CIMT measurements, which may also be high in this study ^[3].

The second point is that there are also large variations in PWMH ^[4]. How great was the variability between different examinations in the same patient, and how great was the inter- and intra-observer variability of the PVWH measurement? It is also important to define the term "periventricular." How was 'periventricular' distinguished from "peripheral," "subcortical," and "deep white matter"?

The third point is that the definition of MCI is arbitrary ^[1]. Five domains were tested, and if abnormal performance was found in two domains or in two tests of a single domain, MCI was diagnosed ^[1]. However, if, for example, two tests in the language domain were impaired, why is it justified to diagnose someone with MCI?

The fourth point is that several different causes of PWMH were not listed as exclusion criteria ^[1]. These include smoking, traumatic brain injury, metabolic encephalopathy, infections, prematurity, demyelinating diseases, cerebral hypoxia, and cardiovascular disease (e.g., atrial fibrillation) ^[1]. If patients with these conditions were included, the results may be misleading.

Finally, we should be informed about how vascular encephalopathy was distinguished from PWMH. PWMH itself may be due to cerebrovascular disease ^[5].

In summary, CIMT and PWMH are poor biomarkers for predicting MIC in patients with Parkinson's disease unless all factors influencing the development of PWMH are included in the analysis.

Declarations**Ethical Approval:** Not applicable.**Consent to Participation:** Not applicable.**Consent for Publication:** Not applicable.**Funding:** None received.**Availability of Data and Material:** All data are available from the corresponding author.**Completing Interests:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.**Author Contribution:** JF was responsible for the design and conception, discussed available data with coauthors, wrote the first draft, and gave final approval. xx and xx: contributed to literature search, discussion, correction, and final approval.**Keywords:** Carotid Intima-Media Wall Thickness, Periventricular White Matter Hyperintensities, Cognitive Decline, Cerebral MRI, Carotid Ultrasound, Parkinson**References**

1. Li S, Che F, Zhang Z. Diagnostic Value of Carotid Intima-Media Thickness Combined with Periventricular White Matter Hyperintensities for Mild Cognitive Impairment in Parkinson's Disease. *Neuropsychiatr Dis Treat*, Nov 24, 2025; 21:2595-2602. Doi: 10.2147/NDT.S553623
2. Meiburger KM, Molinari F, Wong J, Aguilar L, Gallo D, Steinman DA, *et al.* Validation of the Carotid Intima-Media Thickness Variability: Can Manual Segmentations Be Trusted as Ground Truth? *Ultrasound Med Biol*, Jul 2016; 42(7):1598-1611. Doi: 10.1016/j.ultrasmedbio.2016.02.004
3. Salonen R, Haapanen A, Salonen JT. Measurement of intima-media thickness of common carotid arteries with high-resolution B-mode ultrasonography: Inter- and intra-observer variability. *Ultrasound Med Biol*. 1991; 17(3):225-230. Doi: 10.1016/0301-5629(91)90043-v
4. De Kort FAS, Vinke EJ, Van Der Lelij EJ, Anblagan D, Bastin ME, Beiser A, *et al.* Cerebral white matter hyperintensity volumes: Normative age- and sex-specific values from 15 population-based cohorts comprising 14,876 individuals. *Neurobiol Aging*, Feb 2025; 146:38-47. Doi: 10.1016/j.neurobiolaging.2024.11.006
5. Sharma R, Sekhon S, Lui F, *et al.* White Matter Lesions. [Updated 2024 Mar 7]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing, Jan 2025. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK562167/>