



Received: 07-12-2025
Accepted: 17-12-2025

International Journal of Advanced Multidisciplinary Research and Studies

ISSN: 2583-049X

Letter to the Editor

Before Recommending Ultrasound-Guided Superficial Cervical Nerve Block for Musculoskeletal Neck Pain, all Non-Invasive Methods should be Attempted

¹ Josef Finsterer, ² Carla A Scorza, ³ Fulvio Scorza

¹ Department of Neurology, Neurology & Neurophysiology Center, Vienna, Austria

^{2,3} Federal University of São Paulo (UNIFESP/EPMP), São Paulo, Brazil

DOI: <https://doi.org/10.62225/2583049X.2025.5.6.5434>

Corresponding Author: Josef Finsterer

Letter to the Editor

We read with interest the article by Tanigawa *et al.* on a case series of five patients who underwent ultrasound-guided superficial cervical nerve block (SFCNB) in the emergency department (ED) for musculoskeletal neck pain ^[1]. In all five patients, neck pain was reduced from VAS 8–10 to VAS 1–5 ^[1]. The study is noteworthy, but some points require further discussion.

The first point is that oral analgesics and topical lidocaine are not sufficient to diagnose treatment-resistant neck pain ^[1]. In order to indicate more invasive measures for pain treatment, it would have been useful to also use intravenous analgesics, muscle relaxants, and, if necessary, antidepressants before recommending SFCNB. Were opiates also administered to treat neck pain? The disadvantage of SPCNB is that it is invasive, which is why all non-invasive therapies should be exhausted before SFCNB is recommended.

The second point is that the diagnostic evaluation for musculoskeletal neck pain and radiculopathy was not described in detail ^[1]. Were all patients subjected to magnetic resonance imaging (MRI) and X-ray examination of the cervical spine? Patient 1 did not undergo imaging of the cervical spine, even though the patient reported a pain intensity of VAS 10 ^[1]. Patient 5 also did not undergo imaging despite VAS 10 neck pain.

The third point is that no long-term follow-up data were reported ^[1]. How long did the five patients report pain relief? How many days or weeks after the SPCNB was the telephone interview conducted? In how many patients did the pain recur and with what intensity?

The fourth point is that no blood test results were reported to determine whether any of the five patients had an acute or chronic infectious or immunological disease. We should also know how many patients had rheumatic disease, were HLAB27-positive, had Chiari malformation, osteochondrosis, spondylosis, spondylarthritis, listhesis, Hirayama disease, or other confounding conditions.

The fifth point is that neither a depression/anxiety scale nor a stress evaluation was performed to assess the extent to which neck pain was attributable to internal or external stress or depression and anxiety.

The sixth point is that the number of patients was too small to draw general conclusions. In addition, the retrospective design of the study allows only limited conclusions to be drawn from the reported data.

Finally, we should know whether patient 5, who was diagnosed with compression of the C5 nerve by MRI of the cervical spine, had or did not suffer from respiratory insufficiency due to weakness of the diaphragm, which is normally innervated by the C3–5 roots ^[2].

Declarations

Ethical Approval: Not applicable.

Consent to Participation: Not applicable.

Consent for Publication: Not applicable.

Funding: None received.

Availability of Data and Material: All data are available from the corresponding author.

Completing Interests: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contribution: JF was responsible for the design and conception, discussed available data with coauthors, wrote the first draft, and gave final approval. xx: contributed to literature search, discussion, correction, and final approval.

Acknowledgements: None.

Keywords: Neck Pain, Radiculopathy, Ultrasound, Analgesics, Superficial Cervical Nerve Block

References

1. Tanigawa M, Kobayashi K, Naraghi L. A Case Series: Ultrasound-Guided Superficial Cervical Plexus Nerve Block for Musculoskeletal Neck Pain. *J Emerg Med*, Mar 20, 2025; 76:1-6. Doi: 10.1016/j.jemermed.2025.03.019
2. Oliver KA, Ashurst JV. Anatomy, Thorax, Phrenic Nerves. [Updated 2023 Jul 24]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing, Jan 2025. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK513325/>