



Received: 17-12-2025

Accepted: 27-01-2026

ISSN: 2583-049X

**Chemical composition of methyl orange determined by IR spectroscopy****Ioana Stanciu**

Department of Physical Chemistry, Faculty of Chemistry, University of Bucharest, 4-12 Elisabeta Blvd, 030018, Bucharest, Romania


Corresponding Author: **Ioana Stanciu****Abstract**

Methyl orange (also known as helianthine or acid Orange 52) is a synthetic azo dye, commonly used as a pH indicator in titrations. Its chemical formula is  $C_{14}H_{14}N_3NaO_3S$ , with a molecular weight of 327.33 g/mol. Its structure includes an azo group (-N=N-) linking two benzene rings, one of which is substituted with a sulfonic group (-SO<sub>3</sub>Na) and the other with a dimethylamino group (-N(CH<sub>3</sub>)<sub>2</sub>).


Methyl orange is soluble in water and exhibits a significant color change depending on pH: it turns red in acidic medium (pH < 3.1) and yellow in basic medium (pH > 4.4), with a color transition between these values. This property makes it useful in acid-base analysis, especially in titrations of weak acids with strong bases.

**Keywords:** Composition, Methyl Orange, Spectroscopy, FTIR**Introduction**

Methyl orange (helianthine or orange-III) is an orange-colored pH indicator, often used in titrations due to its easily observable color change. It has the molecular formula  $C_{14}H_{14}N_3NaO_3S$  and a molecular mass of 327.33 g/mol. It is an azo dye, being the sodium salt of 4-[4-(dimethylamino)phenylazo]benzenesulfonic acid. Because it changes color under the action of a medium-strength acid, it is used especially for the purpose of titrating acids. Unlike a universal indicator, methyl orange does not show the full color spectrum when changing color. Methyl orange is red in acidic medium and yellow in basic medium. To obtain methyl orange, sulfanilic acid is diazotized with nitric acid and thus converted into 4-benzene diazonium sulfonic acid [1-7].



Subsequently, 4-benzene diazonium sulfonic acid is coupled with N,N-dimethylaniline and treated with sodium hydroxide, transforming into the sodium salt.



Methyl orange is used primarily as an indicator because of the visible color change that occurs. Thus, when used in a solution that is increasingly acidic, the color will change from yellow to orange and finally to red in the most acidic solution. If the process is reversed, the color will change again, from red to orange and finally to yellow in the alkaline solution. Methyl orange has a pK<sub>a</sub> in water of 3.47 at 25 °C [8-12].

## Materials and methods

Measurements were carried out using Bruker Alpha Spectrophotometer, utilizing the ATR (Attenuated Total Reflection) FT-IR sampling technique.



Fig 1: Bruker Alpha Spectrophotometer

## Results and discussions




Fig 2: Spectral indicatorului metil orange

The FTIR spectra of control Methyl orange (Fig 2), displays peak at  $2,924.06 \text{ cm}^{-1}$  for asymmetric  $\text{CH}_3$  stretching vibrations; peaks at  $1,519.78$  and  $1,421.71 \text{ cm}^{-1}$  for the  $\text{C}=\text{C}-\text{H}$  in plane  $\text{C}-\text{H}$  bend; peaks at  $1,040.00$ ,  $1,007.29$ , and  $846.66 \text{ cm}^{-1}$  for ring vibrations; and a peak at  $816.67 \text{ cm}^{-1}$  for the 1, 4 disubstituted (Para) benzene ring<sup>[13-17]</sup>.

## Conclusions

Methyl orange is a synthetic azo dye, with the chemical formula  $\text{C}_{14}\text{H}_{14}\text{N}_3\text{NaO}_3\text{S}$ , which contains an azo group ( $-\text{N}=\text{N}-$ ) linking two benzene rings.

Its structure includes functional groups important for its properties: sulfonic group ( $-\text{SO}_3\text{Na}$ ) and dimethylamino group ( $-\text{N}(\text{CH}_3)_2$ ).

It is soluble in water and acts as a pH indicator, changing color from red in acidic medium to yellow in basic medium.

Its color change property makes it a useful tool in acid-base titrations and chemical analysis.

Its chemical stability and reactivity make it valuable in laboratories and various industrial applications, but it must be used with caution due to its synthetic chemical nature.

## References

- Subagio A, Morita N. Food Chemistry. 2003; 81:97-102.
- Dupont J, White PJ, Carpenter MP, Schaefer EJ, Meydani SN, Elson CE, *et al.* Journal of the American College of Nutrition. 1990; 9(5):438-470.
- Veljković VB, Biberdžić MO, Banković-Ilić IB, Djalović IG, Tasi MB, Nježić ZB, *et al.* Renewable and Sustainable Energy Reviews. 2018; 91:531-548.
- Beadle JB, Just DE, Morgan RE, Reiners RA. Journal of the American Oil Chemists' Society. 1965; 42(2):90-95.
- Strocchi A. Journal of Food Science. 1982; 47(1):36-39.
- Stanciu I. Rheological behaviour of biodegradable lubricant, Journal of Science and Arts. 2019; 3(48):703-708.
- Stanciu I. Rheological investigation of soybean oil from soya beans, Journal of Science and Arts. 2019; 4(49):938-988
- Stanciu I. Modeling the temperature dependence of dynamic viscosity for rapeseed oil, Journal of Science and Arts. 2011; 1:55-58.
- Meneghetti SMP, Meneghetti MR, Wolf CR, Silva EC, Lima GE, Coimbra MDA, *et al.* Journal of the American Oil Chemists' Society. 2006; 83(9):819-822.
- Stanciu I. Journal of Science and Arts. 2018; 18(2):453-458.
- Sheibani A, Ghotbaddini-Bahraman NASER, Sadeghi, FATEMEH. Oriental Journal of Chemistry. 2014; 30(3):1205-1209.
- Stanciu I. Some methods for determining the viscosity index of hydraulic oil, Indian Journal of Science & Technology. 2023; 16(4):254-258.
- Stanciu I. Rheological behavior of corn oil at different viscosity and shear rate, Oriental Journal of Chemistry. 2023; 39(2):335-339.
- Stanciu I. Rheological characteristics of corn oil used in biodegradable lubricant, Oriental Journal of Chemistry. 2023; 39(3):592-595.
- Stanciu I. Effect of temperature on rheology of corn (Zea mays) oil, Oriental Journal of Chemistry. 2023; 39(4):1068-1070.
- Stanciu I. Study Rheological Behavior of Rapeseed oils Compared to Mineral oil, Oriental Journal of Chemistry. 2021; 37(1):247-249.
- Stanciu I. Influence of Temperature on the Rheological Behavior of Orange Honey, Oriental Journal of Chemistry. 2021; 37(2):440-443.