



Received: 20-01-2026
Accepted: 01-03-2026

International Journal of Advanced Multidisciplinary Research and Studies

ISSN: 2583-049X

On \mathbb{Z}_p - linearity

¹ Pombo Jr Dinamérico P, ² Farias Marcelo F, ³ Mauro Patricia CG

¹ Instituto de Matemática e Estatística da UFF, Brazil

² Instituto Multidisciplinar da UFRRJ, Brazil

³ Centro Interdisciplinar de Ciências da Natureza da UNILA, Brazil

Corresponding Author: Pombo Jr Dinamérico P

Abstract

A recent result, valid in the context of certain topological \mathbb{Z}_p - modules vector spaces, is established in the context of topological \mathbb{Z}_p

Keywords: Ring \mathbb{Z}_p of p - Adic Integers, Topological \mathbb{Z}_p - Modules, \mathbb{Z}_p - Linear Mappings

2020 Mathematics Subject Classification: 46H25, 46-03

1. Introduction

In the celebrated paper [2] of Banach it is implicit that, for real normed spaces E and F and for each continuous mapping u from E into F , the conditions

$$u(x + y) = u(x) + u(y) \text{ for all } x, y \in E \text{ (additivity)}$$

and

$$u(\lambda x + \mu y) = \lambda u(x) + \mu u(y) \text{ for all } \lambda, \mu \in \mathbb{R} \text{ and for all } x, y \in E \text{ (linearity)}$$

are equivalent. In [4], the above-mentioned result has been extended to the context of topological vector spaces over certain topological fields. The purpose of the present note is to establish a similar result in the framework of topological \mathbb{Z}_p - modules.

2. Preliminaries and examples

For each prime natural number p , let \mathbb{Z}_p be the discrete valuation ring of p - adic integers [1, 5]; $p\mathbb{Z}_p \supset p^2\mathbb{Z}_p \supset \dots \supset p^n\mathbb{Z}_p \supset p^{n+1}\mathbb{Z}_p \supset \dots$ is a fundamental system of neighborhoods of 0 in \mathbb{Z}_p consisting of ideals of \mathbb{Z}_p , $p\mathbb{Z}_p$ being a maximal ideal. A unitary \mathbb{Z}_p - module E endowed with a topology is a topological \mathbb{Z}_p - module [3, 6] if the mappings $(x, y) \in E \times E \mapsto x + y \in E$ and $(\lambda, x) \in \mathbb{Z}_p \times E \mapsto \lambda x \in E$ are continuous ($E \times E$ and $\mathbb{Z}_p \times E$ endowed with the corresponding product topology).

Let us mention a few examples of topological \mathbb{Z}_p - modules.

2.1 For each integer $n \geq 1$, the \mathbb{Z}_p - module $(\mathbb{Z}_p)^n$, endowed with the product topology, is a Hausdorff topological \mathbb{Z}_p - module.

2.2 The \mathbb{Z}_p - module $(\mathbb{Z}_p)^\mathbb{N}$, endowed with the product topology, is a Hausdorff topological \mathbb{Z}_p - module (obviously, 2.1 follows from 2.2).

2.3 For each non-empty set X and for each topological \mathbb{Z}_p - module F , the \mathbb{Z}_p - module $\mathcal{F}(X; F)$ of all mappings from X into F , endowed with the topology of uniform convergence on the finite subsets of X , is a topological \mathbb{Z}_p - module, which is a Hausdorff space if F is a Hausdorff space (obviously, 2.2 follows from 2.3).

2.4 For each non-empty topological space X and for each topological \mathbb{Z}_p -module F , the \mathbb{Z}_p -module $\mathcal{G}(X; F)$ of all continuous mappings from X into F , endowed with the topology of uniform convergence on the compact subsets of X , is a topological \mathbb{Z}_p -module, which is a Hausdorff space if F is a Hausdorff space.

2.5 For arbitrary topological \mathbb{Z}_p -modules E and F , the \mathbb{Z}_p -module $\mathcal{L}(E; F)$ of all continuous \mathbb{Z}_p -linear mappings from E into F , endowed with the topology of uniform convergence on the bounded subsets of E [5, p. 114], is a topological \mathbb{Z}_p -module, which is a Hausdorff space if F is a Hausdorff space.

3. The result

We shall prove the following:

Proposition 3.1. Let E and F be the topological \mathbb{Z}_p -modules, F being a Hausdorff space. For a mapping $u: E \rightarrow F$, the following conditions are equivalent:

- $u(x + y) = u(x) + u(y)$ for all $x, y \in E$, and u is continuous at $0 \in E$;
- $u(\lambda x + \mu y) = \lambda u(x) + \mu u(y)$ for all $\lambda, \mu \in \mathbb{Z}_p$ and for all $x, y \in E$, and u is uniformly continuous.

Proof. It is obvious that (b) implies (a). Thus it remains to show that (a) implies (b). Indeed, since

$$u(x) - u(y) = u(x) + u(-y) = u(x - y)$$

for all $x, y \in E$, the continuity of u at the origin implies its uniform continuity.

Now, let $\lambda \in \mathbb{Z}_p$ and $x \in E$ be arbitrary. We claim that $u(\lambda x) = \lambda u(x)$. In fact, by the additivity of u and the Principle of Finite Induction,

$$u(nx) = nu(x)$$

for all $n \in \mathbb{N}$. Hence, if $n \in \mathbb{Z}$ and $n < 0$,

$$u(nx) = u(-((-n)x)) = -u((-n)x) = (-(-n))u(x) = nu(x).$$

Finally, by the density of \mathbb{Z} in \mathbb{Z}_p ([1], Proposition 1.2.3), there is a sequence $(\lambda_n)_{n \in \mathbb{N}}$ in \mathbb{Z} converging to λ in \mathbb{Z}_p . Since $(\lambda_n x)_{n \in \mathbb{N}}$ converges to λx in E , the continuity of u at λx implies that $(u(\lambda_n x))_{n \in \mathbb{N}}$ converges to $u(\lambda x)$ in F . On the other hand, $u(\lambda_n x) = \lambda_n u(x)$ for all $n \in \mathbb{N}$, and therefore $(u(\lambda_n x))_{n \in \mathbb{N}}$ converges to $\lambda u(x)$ in F . Consequently the assumption that F is a Hausdorff space furnishes $u(\lambda x) = \lambda u(x)$, which concludes the proof.

4. Conclusion

In this short communication it is shown that, under quite general assumptions, \mathbb{Z}_p -linearity comes from additivity.

5. References

- Amice Y. Les nombres p -adiques. Presses Universitaires de France, 1975.
- Banach S. Sur les fonctionnelles linéaires I et II. Studia Math. 1929; 1:211-216 et 223-239.
- Bourbaki N. Commutative Algebra. Hermann, Paris and Addison-Wesley, Reading, Massachusetts, 1972.
- Pombo Jr DP, Farias MF. A note on linearity. Far East J. Math. Education. 2026; 28:45-49.

- Serre J-P. Corps Locaux. Quatrième édition. Hermann, Paris, Actualités Scientifiques et Industrielles 1296, 1968.
- Warner S. Topological Fields. North-Holland, Amsterdam, Notas de Matemática 126, 1989.