



Received: 26-02-2026
Accepted: 06-04-2026

ISSN: 2583-049X

Smart Decentralized Waste Management and Composting Systems

¹ Salifyanji Sinyinza, ² Jessy Nkhata Kapasa

¹ Department of Architecture, Information and Communications University Zambia, Lusaka, Zambia

² Arch., Department of Architecture, Information and Communications University Zambia, Lusaka, Zambia

DOI: <https://doi.org/10.62225/2583049X.2026.6.2.6123>

Corresponding Author: **Salifyanji Sinyinza**

Abstract

Rapid urbanization in Lusaka has sharply increased municipal waste generation, placing significant strain on the city's landfill-oriented collection system. Chunga Landfill, the primary disposal site, now operates beyond its designed capacity, leading to open dumping, frequent fires, drainage blockages, and leachate contamination.

These challenges exacerbate urban flooding, degrade air quality, and contribute to recurrent cholera and other public-health outbreaks. Similar issues have been observed in cities such as Gwalior, India, where inefficient source segregation, long transport distances, and reliance on centralized disposal systems reduce the efficiency of municipal waste management.

Despite the severity of these problems, there is a paucity of research in the Zambian context investigating alternative, decentralized approaches that can complement existing centralized infrastructure.

This study aims to investigate the feasibility and

effectiveness of decentralized waste management solutions in Lusaka, focusing on the potential for community-level segregation stations, small-scale composters, and sensor-enabled collection points to improve waste-handling efficiency and reduce environmental and health risks. Using a mixed-method approach, the research integrates expert interviews with quantitative analysis of Lusaka's waste composition to evaluate the suitability, scalability, and potential impact of localized processing systems.

Findings indicate that decentralized composting could divert 40–55% of organic waste from Chunga Landfill, reduce collection frequency, lower greenhouse gas emissions, and mitigate public-health hazards.

By addressing the gap in knowledge regarding localized waste management interventions in Zambia, this study contributes practical insights for sustainable urban waste governance and policy development.

Keywords: Centralized Waste Management, Decentralized Systems, Source Segregation, Smart Waste Monitoring, Lusaka Waste Infrastructure, Urban Sustainability

1. Introduction

Waste management in rapidly growing cities such as Lusaka, Nairobi, Accra, and Dar-es-Salaam faces persistent challenges—unplanned settlements, limited municipal resources, insufficient landfill capacity, and a lack of household-level waste segregation. In Lusaka, over 850 tons of waste are generated daily, yet only 40–45% is formally collected. The rest accumulates in drains, open spaces, and informal dumpsites, contributing to flooding, disease outbreaks, greenhouse gas emissions, and land degradation.

A key contributor to the inefficiencies is the centralized waste management model, where all waste—regardless of type—must be transported to distant landfills such as Chunga Landfill. This causes:

- Long transport distances and high fuel costs
- Inefficient collection routes
- Overreliance on municipal trucks
- Poor recycling and organic waste recovery
- Increased landfill pressure and methane emissions

Organic waste (food, yard waste, market waste) constitutes 55–65% of household waste in Lusaka, making it the largest and most problematic fraction.

1.1 Problem Statement

Lusaka’s landfill-oriented waste management approach is increasingly unsustainable due to inadequate waste segregation, inefficient collection systems, high transportation costs, and low recycling and composting rates. These challenges contribute to:

1. Overloaded landfill sites
2. Severe environmental degradation
3. Increased methane emissions
4. Blocked drainage and urban flooding
5. High municipal expenditure

Weak community participation in waste reduction initiatives. There is an urgent need for a decentralized, technology-enhanced model capable of improving efficiency, reducing landfill dependency, and supporting sustainable waste processing.

1.2 General Objective

To evaluate and develop a smart decentralized waste management and composting system that enhances waste collection efficiency, reduces landfill dependency, and promotes sustainable resource recovery in urban communities.

1.3 Specific Objectives

1. To assess the current inefficiencies in Lusaka’s centralized waste management system and identify areas suitable for decentralization.
2. To design and evaluate a pilot smart decentralized waste management model integrating IoT monitoring and community composting hubs.
3. To analyze the environmental, economic, and social impacts of adopting a smart decentralized waste management and composting system.

1.4 Research Questions

1. How can a smart decentralized waste management system improve waste collection efficiency in urban communities?
2. To what extent can decentralized composting reduce the volume of organic waste transported to landfills?
3. What environmental, economic, and social benefits can be achieved through integrating IoT-enabled waste monitoring with community composting hubs?

1.5 Justification of the Study

Municipal waste generation is increasing faster than city infrastructure can support. A smart decentralized model introduces resource efficiency, reduces greenhouse gas emissions, lowers collection and transportation costs, and supports Zambia’s Vision 2030 goals on sustainable cities. The study also promotes practical solutions aligned with SDGs 11 and 12.

2. Literature Review

2.1 Waste Management in Developing Countries

Studies in Sub-Saharan Africa show that urban waste management systems are hindered by low funding, poor infrastructure, and inadequate technologies (UN-Habitat, 2021). Centralized systems rely on hauling all waste to landfills, often far from generation points—leading to inefficiencies.

2.2 Rise of Decentralized Waste Management

Decentralized systems process waste closer to the source, reducing transport demand. They typically involve:

- Community composting facilities
- Biogas digesters
- Waste sorting centers
- Recycling hubs

Research shows decentralized systems reduce collection costs by 20–40% and extend landfill life (Gupta & Yadav, 2020).

2.3 Organic Waste and Composting Technologies

Organic waste forms the majority of waste in African cities. Composting—both aerobic and anaerobic—has proven effective for converting organics into useful products. Decentralized composting reduces methane emissions and supports urban agriculture through nutrient-rich compost.

SWM Step / Stage	Process / Technology Used in Lusaka (or current practice)
Waste generation & storage at source	Households, businesses and institutions store their waste in bins, plastic bags, sacks, drums or containers
Set-out for collection	Waste generators place waste at the entrance or curb outside premises on designated collection day.
Primary collection	In peri-urban/unplanned areas, community-based enterprises (CBEs) handle “primary collection.”
Formal / Secondary collection & transport	For planned / conventional areas, waste is collected by either: – Lusaka City Council (LCC) through its Waste Management Unit (WMU)
Fee/payment & registration system	Waste collection/disposal is regulated — generators are supposed to pay fees for waste disposal/collection under LCC/Waste Management regulations.
Final transport to disposal site (landfill)	Collected waste is transported directly to the official landfill, no intermediate transfer or sorting stations for most waste streams.
Final disposal at landfill	The designated site is Chunga Landfill — the main dumpsite for municipal solid waste in Lusaka. Waste is weighed at the landfill using a weighbridge; disposal fees are paid.
Regulation, oversight and sanitation enforcement	The WMU of LCC supervises and enforces waste-management by-laws, regulates who may collect/dispose waste, and contracts private companies or CBEs where applicable.
Public-awareness & community engagement (partial / periodic)	Monthly clean-ups, community campaigns especially in informal settlements, coordination with ward <u>councillors</u> for drainage cleaning and litter removal — often tied to broader city-cleanliness / climate-resilience initiatives.
Current limitations (mixed waste, no segregation or resource recovery)	Waste is generally collected in mixed/unsegregated form; there is no systematic source segregation, no formal sorting or materials-recovery facilities, and no large-scale composting or recycling system in operation citywide.

2.4 Smart Waste Management Technologies (IoT)

Smart waste technologies involve:

- IoT fill-level sensors in bins
- GPS-enabled route optimization
- Weight-sensing smart bins
- Digital payment and reporting platforms
- Data dashboards for municipalities

Research shows IoT systems reduce:

- Truck fuel use by up to 35%
- Overflow incidents by 80%
- Unnecessary collection trips by 45%

2.5 Integration of Smart Technologies with Decentralized Composting

Emerging studies demonstrate synergistic benefits when IoT tools are deployed alongside community composting hubs:

- Sensors detect organic waste accumulation.
- Digital platforms schedule pickup only when needed.
- Data supports predicting compost output.
- Community adoption improves due to transparency.
- This research explores applying such integrated systems in the Lusaka urban context.
- Environmental degradation and public health risks.

3. Methodology

This study adopted a mixed-method research design, integrating qualitative, quantitative, spatial analytical, and technological pilot-testing approaches. Mixed-methods allow for robust triangulation of findings and are widely recommended in urban environmental research for producing comprehensive and context-sensitive insights (Creswell & Plano Clark, 2018) [25]. The methodology therefore combined field surveys, expert interviews, GIS spatial analysis, secondary data review, and a prototype smart waste system evaluation.

3.1 Study Area

The research was conducted in an kalingalinga within Lusaka characterized by medium-density residential clusters, mixed land uses, and moderate-to-high organic waste generation. The area includes approximately:

- 500 households.
- Local markets and food vendors.
- Schools and institutional facilities.
- This context represents typical waste-generation patterns in Lusaka’s growing residential zones.

3.2 Qualitative Component

3.2.1 Semi-Structured Interviews

Semi-structured interviews were conducted with:

- Architects
- Waste management professionals
- Community leaders
- Municipal planners from Lusaka City Council

Semi-structured interviews support flexible and in-depth exploration of expert perspectives (Kvale & Brinkmann, 2015) [26]. They were designed to capture insights on:

1. Gaps in existing waste management systems
2. Community perceptions of decentralized systems
3. Policy and institutional challenges
4. Technological readiness for smart waste solutions

Indicator / Statistic	Value / Comment for Lusaka
Coordinates (lat. lon)	≈ 15°30' S, 28°10' E
Average / Typical Elevation	~ 1,280 m above sea level
Municipal / District Area	~ 360 km ² (Lusaka city district)
Municipality “planning area” / “special area” / Greater-city area	According to a planning-study, a “study area” covering city + adjoining districts is 850 km ² ; but this includes satellite districts (not purely Lusaka city)
Number of Wards / Administrative subdivisions	38 wards under the city council (within 7 constituencies)
Sanitary zones / Sanitation-zone breakdown	I found no public source that provides a consistent, city-wide “number of sanitary zones” for Lusaka. Recent sanitation-data reports highlight that data gaps exist.
Number of Households	No recent official published total for Lusaka city households was found. 2010 nationwide census data give household numbers by province but not a reliable city-level total.
Total Population (district / city)	2,212,301 (2022 census for Lusaka District)

3.2.2 Data Analysis

Qualitative data were analyzed through thematic analysis, following Braun and Clarke’s (2006) [24] six-step coding framework. Recurring themes identified included:

- Barriers to waste segregation
- Policy limitations
- Urban design constraints
- Community willingness to adopt smart technologies
- Opportunities for localized composting

3.3 Quantitative Component

3.3.1 Waste Data Collection

Quantitative data were obtained from:

- Lusaka City Council (LCC) waste management reports
- Zambia Statistics Agency (ZamStats) datasets
- World Bank global waste databases (2020)
- Field household surveys (n = 50)
- On-site waste characterization studies conducted over 7 days

The Datasets Included:

- Waste generation rates
- Waste composition profiles
- Collection efficiency levels
- Household behavior patterns

City	Population (approx. x., 2025)	Waste generated (tonnes/day)	Waste collected (tonnes/day or % of generated)	Implied per-capita waste generation (kg/person/day)
Lusaka	3,470,870	~ 826–1,200 t/day	~ 30%	~ 0.50–0.52 kg/person/day
Ndola	628,595	No reliable public data found	No data	No reliable public data found
Livingstone	177,393	For similar sized municipalities : ~ 0.608 kg/person/day	Not specified	~ 0.608 kg/person/day (from that study)

3.3.2 Comparative Case Study Assessment

A comparative study analyzed decentralized waste systems such as:

- Eco-block models in China
- Community-based composting hubs in Kenya
- Neighborhood-scale initiatives in South Africa

Comparative insights were used to evaluate the adaptability of global decentralized waste models to Lusaka (Yin, 2018) [29].

3.3.3 Waste Diversion Modelling

Organic waste diversion potential was estimated using:

- Standard composting conversion rates
- UNEP urban composting benchmarks (2020)
- Measured organic fractions from field sampling

This modelling allowed projections of environmental benefits, such as:

- Reduced landfill volumes
- Lower methane emissions
- Potential compost output

3.4 Smart Technology and Pilot System Evaluation

A prototype smart decentralized composting and IoT waste monitoring system was deployed to test its operational performance. The pilot system included:

- Three sensor-enabled smart organic waste bins
- Fill-level detectors (ultrasonic sensors)
- Moisture and temperature sensors for compost monitoring
- A 1-ton/month aerobic composting unit
- A real-time data dashboard for monitoring waste levels

Sensor data allowed evaluation of:

- Collection frequency optimization
- Overflow reduction
- Waste accumulation patterns
- Composting efficiency and operational parameters

This real-world testing provided empirical evidence on the feasibility of integrating IoT with decentralized waste processing.

3.5 Spatial Analysis Component

Spatial analysis was performed using GIS tools to map waste generation patterns, optimal hub placements, and accessibility conditions. GIS analysis supports evidence-based waste planning, site selection, and spatial equity assessment (Goodchild, 2007) [30].

3.5.1 Datasets Used

Spatial datasets included:

- Land use and zoning maps (LCC)
- Road networks for analyzing transfer distances
- Population density layers
- Satellite imagery
- Locations of existing waste collection points
- Informal dumpsite coordinates

3.5.2 GIS Analysis Techniques

Key GIS methods used:

- Suitability analysis to identify optimal sites for decentralized composting hubs.
- Buffer mapping to assess household accessibility.
- Network analysis to model efficient waste collection routes.
- Hotspot analysis to identify high-generation waste zones.

The spatial analysis ensured that proposed smart waste hubs were:

- Strategically located
- Accessible to households
- Minimizing transport burdens
- Situated away from sensitive land uses
- Optimally placed to capture the largest waste streams

3.6 Integration and Triangulation of Data

All qualitative, quantitative, spatial, and sensor-generated data were triangulated to generate holistic insights. This strengthened:

- Validation of findings
- Cross-verification of waste volumes
- Identification of community behavior patterns
- Evaluation of system performance
- Environmental impact analysis

4. Data Presentation and Analysis

This chapter presents the findings from the qualitative interviews, quantitative field surveys, IoT pilot testing, and spatial GIS analysis. The results are organized according to the research objectives and show how a smart decentralized waste management and composting system can improve waste collection efficiency, reduce landfill dependency, and enhance resource recovery.

4.1 Demographic and Household Waste Profile Analysis

A total of 50 households were surveyed within the study area.

Table 4.1: Household Demographics

Variable	Percentage (%)
Household Size (4–6 persons)	58
Household Size (2–3 persons)	32
Household Size (7+ persons)	10
Income Level (Low)	46
Income Level (Middle)	42
Income Level (High)	12

Key Findings

- The majority (58%) of households have 4–6 people, consistent with Lusaka’s average.
- Lower-income households (46%) generate a significant amount of organic waste, mainly food scraps.

4.2 Waste Composition Analysis

Seven-day waste characterization produced the following results.

Table 4.2: Waste Composition (kg/week)

Waste Type	Average kg/household/week	Percentage (%)
Organic Waste	7.1	63
Plastics	2.2	20
Paper/Cardboard	0.9	8
Metals & Glass	0.5	4
Others	0.6	5

Analysis

Organic waste forms the largest fraction (63%), confirming suitability for decentralized composting.

Recyclable categories (plastics + paper + metals) form 32%, yet most remain unsegregated.

The high organic content indicates potential to divert waste from landfills by up to 60% through composting hubs.

4.3 Community Behavior and Perceptions

Table 4.3: Household Waste Practices

Behavior	Percentage (%)
Segregate organic waste	18
Segregate plastics/paper	36
Use formal waste collection	52
Dump waste in informal sites	28
Open burning	20

Analysis

Segregation of organic waste is extremely low (18%), primarily due to:

- Lack of bins
- Poor awareness
- Absence of incentives

Informal dumping and open burning remain significant (48% combined), contributing to pollution.

Households expressed 87% willingness to participate in decentralized composting if bins and collection points are provided.

4.4 Qualitative Interview Themes

Interviews with LCC officers, architects, waste experts, and community leaders revealed 5 main themes:

Inefficiency of Centralized Waste Collection Stakeholders reported:

1. Overreliance on Chunga Landfill
 2. Frequent breakdown of collection trucks
 3. Long routes reducing service frequency
- Poor Urban Waste Segregation Culture

Experts highlighted:

- No policy enforcement
- Lack of infrastructure
- Absence of community awareness
- Environmental Degradation

Municipal planners noted:

- Drain blockages
- Increased flooding
- Uncontrolled dumpsites
- Demand for Community-Level Solutions

Community leaders expressed strong support for:

- Local composting hubs
- Smart bins
- Youth-led waste enterprises
- Technological Readiness

Experts noted growing acceptance of:

- IoT waste monitoring
- Digital platforms for waste alerts
- Sensor-based sorting innovations

4.5 Spatial Analysis Results (GIS)

Spatial modelling included suitability mapping, density analysis, and buffer analysis.

Key Spatial Findings

1. Waste generation hotspots corresponded with:
 - High-density residential clusters
 - Market zones
 - School vicinities
2. Ideal composting hub locations were within:
 - 150–300m buffers
 - Areas with road accessibility
 - Zones away from schools and clinics


3. Optimal smart bin cluster sites were identified near:
 - Busy pedestrian paths
 - Shops
 - Bus stops
4. Reductions in transportation distances ranged from:
 - 2.8 km → 0.5 km (82% reduction) for organic waste.
 - This significantly cuts fuel and collection time.

4.6 Smart IoT System Pilot Results

Three smart sensor bins were installed in the study area for 30 days.

Table 4.4: IoT Pilot Performance Metrics

Parameter	Result
Avg. fill-level detection accuracy	92%
Reduction in overflow incidents	74%
Reduction in unnecessary collection trips	41%
Organic waste captured	164 kg/month
Compost output	51 kg/month



Analysis

High fill-level detection accuracy confirms reliability of the IoT sensors.

Overflow incidents reduced by 74%, improving cleanliness. Collection was done based on real-time data, minimizing wasted fuel and time.

4.7 Quantitative Waste Diversion Modelling

Using UNEP composting benchmarks:
 Organic waste diversion potential = 63%
 Of this, composting efficiency = 62–68%
 Landfill reduction potential = 39–48%
 Projected Annual Impact (per 500 households)

Parameter	Value
Landfill waste reduced	128 tons/year
Compost produced	38–45 tons/year
Methane emissions avoided	22–30 tons CO ₂ e/year

4.8 Integrated Analysis (Answering Research Questions)

RQ1: How can a smart decentralized system improve collection efficiency?

Optimized routes cut trips by 41%.

Overflow incidents fell 74%, improving hygiene.

Smart bins provide real-time waste level data.

RQ2: To what extent can decentralized composting reduce landfill waste?

Up to 48% of total waste diverted from landfill.

Organic fraction alone reduces landfill pressure significantly.

Composting hubs generate usable fertilizer.

RQ3: What environmental, economic, and social benefits result?

Environmental Benefits

Reduced methane emissions

Cleaner streets, fewer illegal dumps

Less burning of waste

Economic Benefits

- Lower municipal fuel and transport costs

- Compost sold to gardens and farms

- Employment opportunities in community-run hubs

- **Social Benefits**

Improved sanitation

High community participation

Enhanced environmental awareness

5. Finding and Discussions

5.1 Current State of Waste Management in Lusaka

Data from Lusaka City Council (LCC, 2022) indicate that the city generates approximately 1,200 tonnes of municipal solid waste per day, of which less than 40% is formally collected. The remainder accumulates in informal dumpsites, drains, and open spaces, contributing to flooding and disease outbreaks. Informal waste pickers contribute significantly to recycling efforts; however, their efficiency is limited by the absence of pre-segregated waste streams, which forces them to hand-sort mixed waste under unsafe and labour-intensive conditions (Kaza *et al.*, 2018) [10].

Waste composition studies further reveal that over 60% of Lusaka's waste is organic, representing a major opportunity for localized composting and resource recovery (UNEP, 2020). However, citywide reliance on the Chunga Landfill places increasing strain on municipal budgets and accelerates environmental degradation.

5.2 Identified Barriers to Effective Waste Management

Interviews conducted with architects, LCC officials, and community representatives highlighted three core barriers:

1. Poor Waste Segregation

Most households mix organic, recyclable, and hazardous waste due to the lack of mandatory segregation policies and limited availability of colour-coded bins. This mirrors challenges reported in other African cities such as Nairobi and Accra (World Bank, 2020) [28].

2. Limited Public Awareness

Respondents emphasized weak public education campaigns on waste reduction and recycling. Communities often lack knowledge regarding composting, responsible disposal, and the environmental consequences of mixed waste.

3. Infrastructural Gaps

Few residential or mixed-use buildings incorporate waste-sorting spaces, composting chambers, or designated recycling points. This reflects a broader architectural gap, where waste management is not integrated into building design (Aye & Widjaya, 2006).

5.3 Benefits of Decentralized Composting and Smart Waste Technologies

Case studies reviewed from Kenya, South Africa, and Barcelona demonstrate clear advantages of decentralized systems:

Decentralized Composting

Research shows that local organic waste processing can reduce landfill-bound waste by up to 50% (Hoornweg & Bhada-Tata, 2012) [9].



Kenya's community eco-villages demonstrated that neighborhood composting facilities:

- Diverted significant organic waste
- Produced affordable fertilizer for urban farming
- Reduced transport costs and reliance on municipal trucks

Smart Waste Technologies

Cities such as Seoul and Barcelona have adopted sensor-enabled bins, leading to:

- 25% reductions in waste collection costs through optimized routing
- Increased accountability and reduced overflowing waste hotspots
- Enhanced resident participation through mobile tracking apps (European Environment Agency, 2019) [7]

Application to Lusaka


If applied to Lusaka's high-density neighborhoods:

- Community composting units could significantly ease pressure on Chunga Landfill
- Sensor-based smart bins could strengthen coordination between formal waste contractors and informal waste pickers
- Local fertilizer production could support urban agriculture in peri-urban areas

For effective implementation, architects must collaborate with planners, engineers, and waste authorities to ensure:

- Waste infrastructure aligns with building codes
- Housing policies encourage decentralized waste processing
- Designs remain functional, aesthetic, and socially acceptable

This approach positions Lusaka's urban blocks as active agents in waste recovery, advancing the shift toward circular urban ecosystems.



6. Acknowledgment

I would like to take this opportunity to express my profound gratitude and deep regards to my supervisor Architect Jessy Nkhata Kapasa for her guidance and motivation throughout the course of this research. And also, to my family and friends for the emotional support throughout the process of putting this paper together, all Glory to God.

7. References

1. UNEP. Africa Waste Management Outlook. United Nations Environment Programme, 2020.
2. World Bank. What a Waste 2.0: A Global Snapshot of Solid Waste Management to 2050. World Bank Group, 2022.
3. Chakrabarti R. Sustainable Architecture and Waste Management: Integrating Environmental Design Strategies. New Delhi: TERI Press, 2015.
4. Gissen D. Sub nature: Architecture's Other Environments. New York: Princeton Architectural Press, 2009.
5. Pomponi F, Moncaster A. Circular economy for the built environment: A research framework. *Journal of Cleaner Production*. 2017; 143:710-718.
6. Ahmed S, Natarajan S. Smart waste management systems: A review. *Waste Management*. 2021; 120:191-203.
7. European Environment Agency. Smart Waste Management Solutions in European Cities. EEA Report, 2019.
8. Gissen D. Subnature: Architecture's Other Environments. New York: Princeton Architectural Press, 2009.
9. Hoornweg D, Bhada-Tata P. What a Waste: A Global Review of Solid Waste Management. World Bank, 2012.
10. Kaza S, Yao L, Bhada-Tata P, Van Woerden F. What a Waste 2.0: A Global Snapshot of Solid Waste Management to 2050. World Bank, 2018.
11. UN-Habitat. Solid Waste Management in the World's Cities. Earthscan, 2010.
12. UNEP. Africa Waste Management Outlook. United Nations Environment Programme, 2020.
13. Blavatnik School of Government. Informal settlements in Lusaka. Oxford. Blavatnik School of Government, 2020.
14. Chiwele D, Lamson-Hall P, Wani S. Informal settlements in Lusaka: Key messages. International Growth Centre / UN-Habitat. International Growth Centre, 2022.
15. ILO / Zambia Statistics Agency. 2022 Census of Population and Housing: Preliminary Results. International Labour Organization, 2022.
16. UN-Habitat. Zambia Urban Housing Sector Profile. UN-Habitat, 2019.
17. Lusaka City Council. About Lusaka. LCC, n.d.
18. Zambia UN Country Team. Common Country Analysis - Zambia. United Nations, 2021.
19. Blavatnik School of Government. Informal settlements in Lusaka. University of Oxford, 2020.
20. International Growth Centre. Informal settlements in Lusaka: Key Messages, 2022.
21. Lusaka City Council. About Lusaka, n.d. <https://www.lcc.gov.zm>
22. UN-Habitat. Zambia Urban Housing Sector Profile, 2019.
23. Zambia Statistics Agency. 2022 Census of Population and Housing: Preliminary Report, 2022.
24. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in Psychology*. 2006; 3(2):77-101.
25. Creswell JW, Plano Clark VL. Designing and conducting mixed methods research (3rd ed.). SAGE Publications, 2018.
26. Kvale S, Brinkmann S. Interviews: Learning the craft of qualitative research interviewing (3rd ed.). SAGE Publications, 2015.
27. UNEP. Global Waste Management Outlook. United Nations Environment Programme, 2020.
28. World Bank. What a Waste 2.0: Global Snapshot of Solid Waste Management, 2020.
29. Yin RK. Case study research and applications: Design and methods (6th ed.). SAGE Publications, 2018.
30. Goodchild MF. Citizens as sensors: The world of volunteered geography. *GeoJournal*. 2007; 69(4):211-221.