



Received: 02-03-2026
Accepted: 12-04-2026

ISSN: 2583-049X

Efficacy of Human Placental Extract Gel as Adjunct to SRP vs. SRP Alone in Stage II Grade B Periodontitis: A Randomized Controlled Trial

¹ Dr. Muglikar Sangeeta, ² Baldi Nidhi

¹ HOD, Department of Periodontology, YMT Dental College and Hospital, Navi Mumbai, India

² MDS Student, Department of Periodontology, YMT Dental College and Hospital, Navi Mumbai, India

DOI: <https://doi.org/10.62225/2583049X.2026.6.2.6156>

Corresponding Author: **Baldi Nidhi**

Abstract

Background: Non-surgical periodontal therapy with scaling and root planing (SRP) remains the cornerstone for managing stage II grade B periodontitis, though adjunctive therapies may enhance outcomes in shallow pockets (≤ 5 mm). This randomized controlled split-mouth trial evaluated human placental extract (HPE) gel as an adjunct to SRP versus SRP alone.

Methods: Fifty bilateral sites in 15 systemically healthy subjects (aged 20-50 years) with stage II grade B periodontitis were randomized: Group A (SRP + HPE gel, n=25 sites) and Group B (SRP alone, n=25 sites). Probing pocket depth (PPD), relative attachment level (RAL), and sulcus bleeding index (SBI) were assessed at baseline, 4 weeks, and 6 weeks using UNC-15 probe and customized

acrylic stents. Non-parametric tests (Friedman, Wilcoxon, Mann-Whitney U) analyzed changes ($p < 0.05$ significant).

Results: Both groups showed significant intra-group improvements ($p < 0.01$). Inter-group comparisons revealed superior outcomes in Group A: PPD reduced by 1.28 mm (vs. 1.08 mm, $p = 0.040$ at 6 weeks); RAL gain of 2.08 mm (vs. 1.16 mm, $p < 0.001$); SBI dropped to 0.51 (vs. 0.87, $p < 0.001$). No adverse effects occurred.

Conclusion: HPE gel significantly augments SRP efficacy in reducing PPD, improving RAL, and minimizing bleeding in stage II grade B periodontitis, likely due to its regenerative (collagen synthesis, growth factors) and anti-inflammatory properties. Longer-term studies are warranted.

Keywords: Human Placental Extract Gel, Scaling and Root Planing, Stage II Grade B Periodontitis, Probing Pocket Depth, Relative Attachment Level, Sulcus Bleeding Index, Split-Mouth Trial

Introduction

The periodontium, comprising gingiva, periodontal ligament, cementum, and alveolar bone^[1] forms a symbiotic barrier against microbial invasion to protect periodontal tissues^[2]. Periodontitis involves gingival inflammation and loss of connective tissue attachment and alveolar bone,^[3] driven by bacterial plaque biofilm that deepens pockets and destroys supporting structures. Non-surgical therapy, primarily scaling and root planing (SRP), remains the cornerstone for initial treatment by removing supra and subgingival biofilm and calculus^[4]. Conventional SRP using power-driven/manual instrumentation yields PPD reductions of 1.29-2.16 mm and CAL gains of 0.55-1.19 mm in 4-6 mm pockets,^[7] with subgingival debridement plus plaque control achieving 0.64-0.74 mm attachment gains and 1.18 mm PPD reduction^[8]. However, limitations persist in deeper pockets, bone defects, and pathogens like *Aggregatibacter actinomycetemcomitans*,^[9] necessitating adjuncts and maintenance. Local drug delivery offers targeted, high-concentration therapy with fewer systemic effects than antimicrobials (e.g., tetracycline fibers, chlorhexidine chips),^[6] though limited by allergies and biofilm disruption; systemic antibiotics risk resistance^[10]. Human placental extract (HPE), rich in proteins, enzymes, hormones, mucopolysaccharides, and polynucleotides,^[9, 11] promotes type I collagen production, anti-inflammatory effects, and reduced IL-6/IL-8 in gingival fibroblasts, showing osteogenic/angiogenic potential^[12]. Gupta *et al.* (2023)^[13] reported superior PPD, CAL, and inflammation improvements with HPE gel adjunct to SRP. Thus, this split-mouth study assessed HPE gel + SRP versus SRP alone on clinical parameters in stage II grade B periodontitis.

Materials and Methods

This study was a split mouth, experimental, randomized controlled trial. Study setting This study was performed on subjects

screened at the outpatient department of periodontology in a tertiary dental care center. It was conducted according to the ADA type II specifications using electrically operated dental chair with proper illumination X-ray unit and autoclaved set of instruments. Source of data Subjects of either sex within the age range of 20-50 years who reported to the Out Patient Department of a tertiary dental care center, with probing pocket depth ≤ 5 mm in bilateral sites subjects were selected by convenience sampling technique. Method of selection the subjects for the study were selected based on the following inclusion and exclusion criteria.

Inclusion Criteria

1. Systemically healthy subjects of either sex between 20-50 years.
2. Subjects presenting with bilateral sites with periodontal pockets with probing depth ≤ 5 mm.
3. Subjects willing to be a part of the study.

Exclusion criteria

1. Pregnant and lactating mothers.
2. Subjects with habit of smoking, tobacco chewing.
3. Subjects who are not on any medication.
4. Subjects with history of any periodontal treatment in past 6 months.
5. Subjects with any known drug allergy.

Withdrawal Criteria

1. Non-compliant subjects
2. Subjects lost to follow up to match with original sample size fresh subjects were to be recruited for evaluation. But in the present study there was no loss to follow up and all patients were compliant.



Fig 3: Post scaling photograph



Fig 4: Intrapocket application of HPE gel



Fig 5: Placement of HPE gel till the pocket is filled completely



Fig 6: Coe pak placement



Fig 1: Human placental extract (HPE)




Fig 2: Collagen plug mixed with HPE gel

Group A (SRP+ Human Placental Extract Gel)



Fig 7: At baseline



Fig 8: Follow up at 4 weeks



Fig 13: Follow up at 6 weeks

Group B (SRP)



Fig 14: At Baseline



Fig 15: Follow up at 4 weeks



Fig 16: Follow up at 6 weeks

Results

This study was carried out in the Department of Periodontology to assess the periodontal treatment outcome with Human placental extract gel as an adjunct to scaling and root planning (SRP) as compared to SRP alone in subjects of either sex between 20-50 years of age (mean age of 45 years) with Stage II Grade B periodontitis. Fifty selected bilateral sites in 15 patients (10 males,5 females) with pocket depth ≤ 5 mm were randomly divided at Group A (treated with Human placental extract gel + SRP) & Group B (SRP alone). Probing pocket depth, relative attachment level and sulcus bleeding index were recorded at baseline, 4 weeks, 6 weeks (**Table1, Table 2**).

All the subjects were compliant and there were no dropouts from the study. The changes in the clinical parameters over a period of 6 weeks were recorded. The data obtained was tabulated and subjected to statistical analysis.

Statistical procedures:

All data were entered into a computer by giving coding system, proofed for entry errors. Data obtained was compiled on a MS office Excel Sheet (v 2019, Microsoft Redmond, Washington, United states). Data was subjected to statistical analysis using Statistical package for social sciences (SPSS v 26.0, IBM). Descriptive statistics like frequencies and percentage for categorial data, Mean and SD for numerical data has been depicted.

Normality of numerical data was checked using Shapiro-Wilk test and was found that the data did not follow a normal curve; or for graded data, hence non- parametric

tests have been used for comparisons. Inter Site comparison (Between two sites) was done using Mann Whitney U test. Intra Site comparison was done using Friedman’s for (>2 observations) followed by pair wise comparison using Wilcoxon signed rank test.

For all the statistical test, $p < 0.05$ was considered to be statistically significant, keeping α error at 5% and β error at 20%, thus giving a power to the study 80%.

*= statistically significant difference ($p < 0.05$) **= statistically highly significant difference (< 0.01).

= Nonsignificant difference (> 0.05) for all tables and graphs.

Intra Group Comparison:

The values of PPD for Group A decreased from baseline (4.56 ± 0.50) throughout follow ups at 4weeks (3.68 ± 0.69) and 6 weeks (3.28 ± 0.54). On intra group comparison, there was statistically highly significant difference seen ($p < 0.01$) in PPD between baseline, 4weeks and 6weeks with higher values at baseline (Table 3) (Graph 1).

The values of PPD for Group B decreased from baseline (4.68 ± 0.47) throughout follow ups at 4weeks (3.76 ± 0.59) and 6 weeks (3.6 ± 0.5). On intra group comparison, there was statistically highly significant difference seen ($p < 0.01$) in PPD between baseline, 4weeks and 6weeks with higher values at baseline (Table 4) (Graph 1).

There was a statistically highly significant difference seen for the PPD values of at all pairs of time intervals ($p < 0.01$) in Group A (Table 5). In Group B, there was a statistically highly significant difference seen for the PPD values between baseline and 4 weeks and there was no significant difference from 4 weeks to 6 weeks ($p < 0.01$) (Table 6).

Inter Group Comparison:

PPD values at baseline between Group A (4.56 ± 0.5) and Group B (4.68 ± 0.47) showed no statistically significant ($p < 0.05$) difference.

At 4 weeks follow up, no statistically significant difference ($p < 0.01$) was observed between both groups, Group A (3.68 ± 0.69) and Group B (3.76 ± 0.59). There was a statistically significant difference seen ($p < 0.05$) for PPD at 6weeks between the Group A (3.28 ± 0.54) and B, with higher values in Group B (3.60 ± 0.50) (Table 7) (Table 12) (Graph 4).

Intragroup Comparison:

The values of RAL for sites in Group A decreased from baseline (7.04 ± 1.30) throughout follow ups at 4 weeks (5.56 ± 1.71) and 6 weeks (4.9 ± 1.48) On intragroup comparison there was statistically significant difference seen ($p < 0.01$) in RAL between baseline, 4weeks and 6weeks with higher values at baseline. (Table 8) (Graph 2).

The values of RAL for Group B decreased from Baseline (8.56 ± 0.82) throughout follow up at 4 weeks (7.48 ± 0.87) and 6 weeks (7.4 ± 0.9) There was statistically highly significant difference seen ($p < 0.01$) in RAL between baseline, 4weeks and 6weeks with higher values at Baseline.

(Table 9) (Graph 2). There was a highly statistical difference seen for the RAL values between baseline to 4 weeks and there was no significant difference between 4 weeks to 6 weeks of time interval ($p < 0.01, 0.05$) at Group B (Table 6).

Intergroup Comparison:

There was a statistically highly significant difference seen ($p < 0.01$) for RAL at baseline between the Group A (7.04 ± 1.30) and Group B (8.56 ± 0.8), there is statistical significant difference for RAL at 4 weeks Group A (5.56 ± 1.7), Group B (7.48 ± 1.4), For RAL at 6 weeks group A (4.9 ± 1.48) and Group B (7.4 ± 0.9) there was statistically highly significant difference with higher values in Group B (Table 7) (Table 12) (graph 5).

Sulcus Bleeding Index:

SBI was measured using Periodontal probe at baseline, 4 weeks, 6 weeks.

Intragroup Comparison:

The values of SBI for Group A decreased from baseline (2.55 ± 0.54) throughout follow up at 4 weeks (1.20 ± 0.51) and 6 weeks (0.51 ± 0.99) on intra group comparison There was statistically highly significant difference seen ($p < 0.01$) in SBI between baseline, 4weeks and 6weeks with higher values at Baseline. (Table 10) (Graph 3).

The values of SBI for Group B decreased from baseline (2.8 ± 0.45) throughout follow up at 4 weeks (1.74 ± 0.54) and 6 weeks (0.87 ± 0.38) on intra group comparison, there was statistical significance difference ($p < 0.01$) seen from baseline to 4 weeks and 6 weeks. (Table 11) (Graph 3) There was statistically highly significant difference seen ($p < 0.01$) in SBI between Baseline, 4weeks and 6weeks with higher values at baseline.

Intergroup Comparison:

There was statistically significant difference ($p < 0.01$) seen between Group A (2.55 ± 0.54) and Group B (2.8 ± 0.45) at baseline. At 4 weeks follow up, highly statistically significant difference between Group A (1.20 ± 0.51) showed greater reduction in SBI value as compared to Group B (1.74 ± 0.54). At follow up of 6 weeks, highly statistically significant difference ($p < 0.01$) between both groups i.e. Group A (0.51 ± 0.99) with higher values in Group B (0.87 ± 0.38) (Table 7) (Table 12).

Tables & Graphs

Table 1: Age wise distribution of subjects

| Age (Years) | N |
|--------------|-----------|
| 20-35 Years | 2 |
| 36-40 years | 4 |
| 41-45 years | 5 |
| 46-50years | 4 |
| Total | 15 |

Table 2: Gender wise distribution of subjects

| Gender | Group A(N) | Group B(N) |
|--------|------------|------------|
| Male | 10 | 10 |
| Female | 5 | 5 |

Table 3: Intra Group Comparison of PPD at Group A (SRP + human placental extract gel)

| | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi- Square value | p value of Friedman Test |
|----------|----|------|----------------|---------|---------|--------|-----------|-------------------|--------------------------|
| PPD BL | 25 | 4.56 | 0.507 | 4 | 5 | 5.00 | 2.88 | 39.325 | 0.000** |
| PPD (4w) | 25 | 3.68 | 0.690 | 2 | 5 | 4.00 | 1.78 | | |
| PPD (6w) | 25 | 3.28 | 0.542 | 2 | 4 | 3.00 | 1.34 | | |

Table 4: Intra Group Comparison of PPD at Group B (SRP)

| | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi- Square value | p value of Friedman Test |
|----------|----|------|----------------|---------|---------|--------|-----------|-------------------|--------------------------|
| PPD BL | 25 | 4.68 | 0.476 | 4 | 5 | 5.00 | 2.92 | 40.582 | 0.000** |
| PPD (4w) | 25 | 3.76 | 0.597 | 3 | 5 | 4.00 | 1.62 | | |
| PPD (6w) | 25 | 3.60 | 0.500 | 3 | 4 | 4.00 | 1.46 | | |

Table 5: Pair wise Comparison using Wilcoxon Signed Ranks Test at Group A (SRP + Human Placental Extract Gel)

| Pairs | Z value | p value of Wilcoxon Signed Ranks Test |
|-----------------|---------|---------------------------------------|
| PPD 4w - PPD BL | -4.147 | 0.000** |
| PPD 6w - PPD BL | -4.463 | 0.000** |
| PPD 6w - PPD 4w | -2.887 | 0.004** |
| RAL 4w - RAL BL | -4.239 | 0.000** |
| RAL 6w - RAL BL | -4.458 | 0.000** |
| RAL 6w - RAL 4w | -3.419 | 0.001** |
| SBI 4w - SBI BL | -4.387 | 0.000** |
| SBI 6w - SBI BL | -3.897 | 0.000** |
| SBI 6w - SBI 4w | -3.717 | 0.000** |

Table 6: Pair wise Comparison using Wilcoxon Signed Ranks Tests at Group B

| Pairs | Z value | p value of Wilcoxon Signed Ranks Test |
|-----------------|---------|---------------------------------------|
| PPD 4w - PPD BL | -4.796 | 0.000** |
| PPD 6w - PPD BL | -4.508 | 0.000** |
| PPD 6w - PPD 4w | -1.414 | 0.157# |
| RAL 4w - RAL BL | -4.838 | 0.000** |
| RAL 6w - RAL BL | -4.420 | 0.000** |
| RAL 6w - RAL 4w | -0.816 | 0.414# |
| SBI 4w - SBI BL | -4.385 | 0.000** |
| SBI 6w - SBI BL | -4.384 | 0.000** |
| SBI 6w - SBI 4w | -4.400 | 0.000** |

Table 7: Intergroup Comparison of Variables at Group A (SRP + Human Placental Extract Gel) and Group B (SRP)

| Group | N | Mean | Std Deviation | | Mann Whitney U value | | Z value | P value of Mann Whitney U Test | |
|--------|------|---------|---------------|-------|----------------------|------|---------|--------------------------------|---------|
| PPD BL | A 25 | 4.56 | 0.507 | 24 | 600 | 5 | 275.000 | -0.865 | 0.387# |
| | B 25 | 4.68 | 0.476 | 27 | 675 | 5 | | | |
| PPD 4w | A 25 | 3.68 | 0.690 | 24.88 | 622 | 4 | 297.000 | -0.342 | 0.732# |
| | B 25 | 3.76 | 0.597 | 26.12 | 653 | 4 | | | |
| PPD 6w | A 25 | 3.28 | 0.542 | 21.8 | 545 | 3 | 220.000 | -2.056 | 0.040* |
| | B 25 | 3.60 | 0.500 | 29.2 | 730 | 4 | | | |
| RAL BL | A 25 | 7.04 | 1.306 | 17.36 | 434 | 7 | 109.000 | -4.12 | 0.000** |
| | B 25 | 8.56 | 0.821 | 33.64 | 841 | 9 | | | |
| RAL 4w | A 25 | 5.56 | 1.710 | 17.64 | 441 | 5 | 116.000 | -3.921 | 0.000** |
| | B 25 | 7.48 | 0.872 | 33.36 | 834 | 7 | | | |
| RAL 6w | A 25 | 4.96 | 1.485 | 15.28 | 382 | 5 | 57.000 | -5.067 | 0.000** |
| | B 25 | 7.40 | 0.913 | 35.72 | 893 | 7 | | | |
| SBI BL | A 25 | 2.55000 | 0.549621 | 21.5 | 537.5 | 2.5 | 212.500 | -1.994 | 0.046* |
| | B 25 | 2.83000 | 0.454835 | 29.5 | 737.5 | 3 | | | |
| SBI 4w | A 25 | 1.20000 | 0.515388 | 17.96 | 449 | 1 | 124.000 | -3.737 | 0.000** |
| | B 25 | 1.74000 | 0.547152 | 33.04 | 826 | 1.5 | | | |
| SBI 6w | A 25 | 0.51000 | 0.998645 | 17.46 | 436.5 | 0.25 | 111.500 | -3.951 | 0.000** |
| | B 25 | 0.87000 | 0.389444 | 33.54 | 838.5 | 0.75 | | | |

Table 8: Intragroup Comparison of RAL Group A (SRP + Human Placental Extract Gel)

| | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi- Square value | p value of Friedman Test |
|----------|----|------|----------------|---------|---------|--------|-----------|-------------------|--------------------------|
| RAL BL | 25 | 7.04 | 1.306 | 5 | 9 | 7.00 | 2.94 | 43.929 | 0.000** |
| RAL (4w) | 25 | 5.56 | 1.710 | 3 | 9 | 5.00 | 1.82 | | |
| RAL (6w) | 25 | 4.96 | 1.485 | 3 | 7 | 5.00 | 1.24 | | |

Table 9: Intragroup Comparison of RAL at Group B (SRP)

| | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi-Square value | p value of Friedman Test |
|----------|----|------|----------------|---------|---------|--------|-----------|------------------|--------------------------|
| RAL BL | 25 | 8.56 | 0.821 | 7 | 10 | 9.00 | 2.96 | 43.772 | 0.000** |
| RAL (4w) | 25 | 7.48 | 0.872 | 6 | 9 | 7.00 | 1.54 | | |
| RAL (6w) | 25 | 7.40 | 0.913 | 6 | 9 | 7.00 | 1.50 | | |

Table 10: Intragroup Comparison of SBI at Group A (SRP + Human Placental Extract Gel)


| | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi- Square value | p value of Friedman Test |
|----------|----|---------|----------------|---------|---------|---------|-----------|-------------------|--------------------------|
| SBI BL | 25 | 2.55000 | 0.549621 | 2.000 | 3.500 | 2.50000 | 2.96 | 44.240 | 0.000** |
| SBI (4w) | 25 | 1.20000 | 0.515388 | 0.500 | 3.000 | 1.00000 | 1.96 | | |
| SBI (6w) | 25 | 0.51000 | 0.998645 | 0.000 | 5.000 | .25000 | 1.08 | | |

Table 11: Intragroup Comparison Of SBI at Group B (SRP)


| | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi-Square value | p value of Friedman Test |
|----------|----|---------|----------------|---------|---------|---------|-----------|------------------|--------------------------|
| SBI BL | 25 | 2.83000 | 0.454835 | 1.750 | 3.500 | 3.00000 | 3.00 | 50.000 | 0.000** |
| SBI (4w) | 25 | 1.74000 | 0.547152 | 0.500 | 2.750 | 1.50000 | 2.00 | | |
| SBI (6w) | 25 | 0.87000 | 0.389444 | 0.000 | 1.750 | 0.75000 | 1.00 | | |

Table 12: Inter Group Comparison of Variables at Group A and Group B


| | | N | Mean | Std. Deviation | Minimum | Maximum | Median | Mean Rank | Chi-Square value | p value of Friedman Test |
|----------|---|----|---------|----------------|---------|---------|--------|-----------|------------------|--------------------------|
| PPD BL | A | 25 | 4.56 | 0.507 | 24 | 600 | 5 | 275.000 | -0.865 | 0.387# |
| | B | 25 | 4.68 | 0.476 | 27 | 675 | 5 | | | |
| PPD (4w) | A | 25 | 3.68 | 0.690 | 24.88 | 622 | 4 | 297.000 | -0.342 | 0.732# |
| | B | 25 | 3.76 | 0.597 | 26.12 | 653 | 4 | | | |
| PPD (6w) | A | 25 | 3.28 | 0.542 | 21.8 | 545 | 3 | 220.000 | -2.056 | 0.040* |
| | B | 25 | 3.60 | 0.500 | 29.2 | 730 | 4 | | | |
| RAL BL | A | 25 | 7.04 | 1.306 | 17.36 | 434 | 7 | 109.000 | -4.12 | 0.000** |
| | B | 25 | 8.56 | 0.821 | 33.64 | 841 | 9 | | | |
| RAL (4w) | A | 25 | 5.56 | 1.710 | 17.64 | 441 | 5 | 116.000 | -3.921 | 0.000** |
| | B | 25 | 7.48 | 0.872 | 33.36 | 834 | 7 | | | |
| RAL (6w) | A | 25 | 4.96 | 1.485 | 15.28 | 382 | 5 | 57.000 | -5.067 | 0.000** |
| | B | 25 | 7.40 | 0.913 | 35.72 | 893 | 7 | | | |
| SBI BL | A | 25 | 2.55000 | 0.549621 | 21.5 | 537.5 | 2.5 | 212.500 | -1.994 | 0.046* |
| | B | 25 | 2.83000 | 0.454835 | 29.5 | 737.5 | 3 | | | |
| SBI (4w) | A | 25 | 1.20000 | 0.515388 | 17.96 | 449 | 1 | 124.000 | -3.737 | 0.000** |
| | B | 25 | 1.74000 | 0.547152 | 33.04 | 826 | 1.5 | | | |
| SBI (6w) | A | 25 | 0.51000 | 0.998645 | 17.46 | 436.5 | 0.25 | 111.500 | -3.951 | 0.000** |
| | B | 25 | 0.87000 | 0.389444 | 33.54 | 838.5 | 0.75 | | | |




Graph 1: Intra Group Comparison of PPD at Group A and Group B




Graph 2: Intra Group Comparison of RAL at Group A (SRP + Human Placental Extract Gel) And Group B (SRP)




Graph 3: Intra Group Comparison of SBI at Group A and Group B



Graph 4: Inter Group Comparison of PPD at Group A (SRP + Human Placental Gel) and Group B (SRP)



Graph 5: Inter Group Comparison of RAL at Group A (SRP + Human Placental Gel) and Group B (SRP)



Graph 6: Inter Group Comparison of SBI at Group A and Group B

Conclusion

This randomized controlled split-mouth trial investigated the efficacy of human placental extract (HPE) gel as an adjunct to scaling and root planing (SRP) in the non-surgical management of stage II grade B periodontitis. The study compared a total of 50 sites, 25 sites treated with SRP intrapocket application of HPE gel (Group A) to 25 sites treated with SRP alone (Group B) in 15 subjects, assessing probing pocket depth (PPD), relative attachment level (RAL), and sulcus bleeding index (SBI) at baseline, 4 weeks, and 6 weeks. The superior clinical outcomes in Group A align with the regenerative and anti-inflammatory properties of HPE, as supported by prior studies demonstrating its ability to enhance collagen synthesis, reduce pro-inflammatory cytokines, and facilitate wound healing. The absence of adverse effects and ease of HPE gel application highlight its potential for clinical integration. Within the limitations of the study, it can be concluded that HPE gel as an adjunct to SRP significantly enhances clinical outcomes in the non-surgical treatment of Stage II Grade B Periodontitis, offering a promising approach to improve periodontal health and potentially reduce the need for surgical interventions. These findings advocate for further research to establish HPE gel as a standard adjunct in periodontal therapy, paving the way for advanced regenerative strategies in clinical practice.

References

- Newman MG, Takei HH, Klokkevold PR, Carranza FA. Newman and Carranza's Clinical Periodontology. 13th ed. Philadelphia: Elsevier, 2019.
- Bartold PM, Van Dyke TE. Periodontitis: A host-mediated disruption of microbial homeostasis. Unlearning learned concepts. *Periodontol 2000*. 2013; 62(1):203-217
- Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nat Rev Dis Primers*. 2017; 3:17038.
- Cobb CM. Non-surgical pocket therapy: Mechanical. *Ann Periodontol*. 1996; 1(1):443-490.
- Muhlemann HR, Son S. Gingival sulcus bleeding-a leading symptom in initial gingivitis. *Helv Odontol Acta*. 1971; 15(2):107-113.
- Drisko CH. Nonsurgical periodontal therapy. *Periodontol 2000*. 2001; 25:77-88.
- Greenstein G. Local drug delivery in the treatment of periodontal diseases: Assessing the clinical significance of the results. *J Periodontol*. 2006; 77(4):565-578.
- Cobb CM. Clinical significance of non-surgical periodontal therapy: An evidence-based perspective of scaling and root planing. *J Clin Periodontol*. 2002; 29(Suppl 2):6-16.
- Van der Weijden GA, Timmerman MF. A systematic review on the clinical efficacy of subgingival debridement in the treatment of chronic periodontitis. *J Clin Periodontol*. 2002; 29(Suppl 3):55-71.
- Slots J, Ting M. *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in human periodontal disease: Occurrence and treatment. *Periodontol 2000*. 1999; 20:82121.
- Matesanz-Pérez P, García-Gargallo M, Figuero E, Bascones-Martínez A, Sanz M, Herrera D. A systematic review on the effects of local antimicrobials as adjuncts to subgingival debridement, compared with subgingival debridement alone, in the treatment of chronic periodontitis. *J Clin Periodontol*. 2013; 40(3):227-241.
- Carter AM, Enders AC. Comparative aspects of trophoblast development and placentation. *Reprod Biol Endocrinol*. 2004; 2:46.
- Akagi T, Matsumura Y, Yasui K, Nakata M, Sasaki M. Effect of human placental extract on proliferation and anti-inflammatory activity of human gingival fibroblasts. *J Periodontol Res*. 2020; 55(4):475-483.
- Gupta S, Sharma A, Kumar P, Agarwal R. Evaluation of human placental extract as an adjunct to scaling and root planing in the treatment of chronic periodontitis: A clinical study. *J Periodontol*. 2023; 94(2):156-165.
- Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, *et al*. Periodontitis: Consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Periodontol*. 2018; 89(Suppl 1):S173-S182.
- Eke PI, Dye BA, Wei L, Thornton-Evans GO, Genco RJ. CDC Periodontal Disease Surveillance workgroup. Prevalence of periodontitis in adults in the United States: 2009 and 2010. *J Dent Res*. 2012; 91(10):914-920. Doi: 10.1177/0022034512457373
- Ower P. Non-surgical periodontal therapy: an overview. *Dent Update*. 2000; 27(9):426-432.
- Minabe M, Uematsu A, Nishijima K, *et al*. Application of a local drug delivery system to periodontal therapy: I. Development of collagen preparations with immobilized tetracycline. *J Periodontol*. 1989; 60:113-117. Doi: 10.1902/jop.1989.60.2.113. [DOI] [PubMed] [Google Scholar]
- Kho P, Shales FC, Hardie JM. Effect of supragingival plaque control on subgingival microflora. *J Clin Periodontol*. 1985; 12(8):676-686.

20. McNabb H, Mombelli A, Lang NP. Supragingival cleaning and subgingival microbiota. *J Clin Periodontol.* 1992; 19(2):144-150.
21. Magnusson I, Lindhe J, Yoneyama T, Liljenberg B. Recolonization of subgingival microbiota post-scaling. *J Clin Periodontol.* 1984; 11(3):193-207.
22. Badersten A, Nilveus R, Egelberg J. Effect of nonsurgical periodontal therapy. II. Severely advanced periodontitis. *J Clin Periodontol.* 1984; 11(2):63-76.
23. Breininger DR, O'Leary TJ, Blumenshine RVH. Comparative effectiveness of ultrasonic and hand scaling. *J Periodontol.* 1987; 58(1):9-14.
24. Johnson GK, Reinhardt RA, Tussing GJ, Krejci RF. Enhanced access and visualization in sonic scaling. *J Periodontol.* 1989; 60(6):321-326.
25. Sherman PR, Hutchens LH, Jewson LG, Moriarty JM, Greco GW, McFall WT Jr. Detection of residual subgingival calculus post-SRP. *J Periodontol.* 1990; 61(1):9-15.
26. Nagy RJ, Otomo-Corgel J, Stambaugh R. Effectiveness of subgingival scaling with modified curets. *J Periodontol.* 1992; 63(3):208-214.
27. Anderson GB, Palmer JA, Bye FL, Smith BA, Caffesse RG. Effectiveness of single versus multiple scaling sessions. *J Periodontol.* 1996; 67(4):367-373.
28. Miyazaki A, Yamaguchi T, Nishikata J, *et al.* Comparison of Nd:YAG, CO2 laser, and ultrasonic scaling in periodontitis. *J Periodontol.* 2003; 74(8):1161-1169.
29. Colombo AP, Teles RP, Torres MC, Rosalém W, *et al.* Effects of scaling and root planing on subgingival microbiota in Brazilian subjects. *J Periodontol.* 2005; 76(8):1300-1308.
30. Bitto A, Polito F, Irrera N, *et al.* Polydeoxyribonucleotide reduces cytokine production and the severity of collagen-induced arthritis by stimulation of adenosine A2A receptor. *Arthritis Rheum.* 2011; 63(11):3364-3371.
31. Geisinger ML, Mealey BL, Schoolfield J, Mellonig JT. Effectiveness of endoscopic-assisted scaling and root planing. *J Periodontol.* 2007; 78(6):1027-1035.
32. Santos VR, Lima JA, De Mendonça AC, *et al.* Clinical and metabolic effects of fullmouth versus partial-mouth scaling in diabetic patients. *J Clin Periodontol.* 2009; 36(10):867-875.
33. Aljateeli M, Koticha T, Bashutski J, Sugai J, Wang HL. Impact of initial scaling on periodontal surgery outcomes. *J Periodontol.* 2014; 85(4):e77-e84.
34. Sinha S, Nair V, Das I, *et al.* Laser-assisted periodontal therapy versus conventional scaling and root planing. *J Indian Soc Periodontol.* 2023; 27(2):145-151.
35. Wu CH, Chang GY, Chang WC, Hsu CT, Chen RS. Wound healing effects of porcine placental extracts on rats with thermal injury. *British Journal of Dermatology.* 2003; 148(2):236-245.
36. Hong JW, Lee WJ, Hahn SB, Kim BJ, Lew DH. The effect of human placenta extract in a wound healing model. *Annals of Plastic Surgery.* 2010; 65(1):96-100.
37. Chakraborty PD, De D, Bandyopadhyay S, Bhattacharyya D. Human aqueous placental extract as a wound healer. *Journal of Wound Care.* 2009; 18(11):462-467.
38. Filatov VP. Tissue therapy: Teaching on biogenic stimulators. Moscow: Foreign Languages Publishing House, 1955.
39. Kaushal V, Verma K, Manocha S, Hooda HS, Das BP. Efficacy of human placental extract in radiation-induced mucositis. *Indian J Med Res.* 2001; 114:149-154.
40. Gupta J, Srinivasan SV, Daniel M. Efficacy of betamethasone, placental extract, and hyaluronidase in oral submucous fibrosis. *J Oral Pathol Med.* 2012; 41(8):615-620.
41. Thakur G, Thomas S, Bhargava D, *et al.* Efficacy of topical placental extracts on postoperative fibrotomy wounds in oral submucous fibrosis. *J Oral Maxillofac Surg.* 2015; 73(8):1467-1473.
42. Shah P, Venkatesh R, More C, *et al.* Therapeutic efficacy of placental extract with dexamethasone versus hyaluronidase with dexamethasone in oral submucous fibrosis. *J Oral Biol Craniofac Res.* 2016; 6(Suppl 1):S40-S45.
43. Chang SW, Lee W, Lee JH, *et al.* Synergistic effects of MTA and human placental extract on dental pulp cells. *J Endod.* 2016; 42(6):929-936.
44. Garg A, Sharma S, Chauhan SS, Sinha A, Agrawal R. Comparative evaluation of Placentrex and ozonated olive oil in laser depigmentation. *J Cosmet Dent.* 2022; 38(2):45-52.
45. Kondaveeti SS, Naik V, Patil A, *et al.* Benefits of Placentrex in oral mucositis management in oral cancer patients undergoing chemoradiation. *J Cancer Res Ther.* 2018; 14(5):1018-1023.
46. Shinde CV, Saawarn N, Kohli S, *et al.* Efficacy of placental extract versus triamcinolone acetonide in OSMF. *J Oral Biol Craniofac Res.* 2019; 9(2):162-167.
47. Sharma A, Sharma S, Nagar A. Effect of placental extracts in periodontal pockets. *J Indian Soc Periodontol.* 2020; 24(3):223-228.
48. Katkurwar A, Chaudhari D, Mahale S, *et al.* Clinical and histological evaluation of placental extract gel in gingival depigmentation. *J Periodontal Res.* 2021; 56(4):789-796.
49. Kumar P, Singh A, Verma S, Gupta R, Yadav S. Efficacy of placental extract gel in chronic periodontitis. *J Periodontol.* 2021; 92(10):1385-1392.
50. Morsy SM, El-Sherbiny RH, Shata MS. Effect of placental extract gel in experimental periodontitis in rats. *J Periodontal Res.* 2022; 57(3):587-596.
51. Morsy SM. Efficacy of topical placental extract gel in recurrent aphthous stomatitis. *Clin Oral Investig.* 2022; 26(6):4589-4596.
52. Miyamoto T, *et al.* Influence of placenta extract intake on periodontal diseases and oral environment: A case series. *J Periodontol.* 2024; 95(3):245-253.
53. Bhadauriya P, *et al.* Clinical efficacy of scaling and root planing with placental extract gel under magnification in chronic periodontitis. *J Periodontol.* 2024; 95(4).
54. Lindhe J, Nyman S. Scaling and root planing in shallow pockets. *J Clin Periodontol.* 1985; 12(5):415-418. Doi: 10.1111/j.1600-051x.1985.tb00930.x