



Received: 29-03-2026  
Accepted: 09-05-2026

## International Journal of Advanced Multidisciplinary Research and Studies

ISSN: 2583-049X

### Proposed $T_{bb}$ Tetraquark at $10.5765 \text{ GeV}/c^2$

JJ Bevelacqua

Bevelacqua Resources, 7531 Flint Crossing Circle SE, Owens Cross Roads, AL 35763, USA

Corresponding Author: JJ Bevelacqua

#### Abstract

The proposed  $T_{bb}$  tetraquark at  $10.5765 \text{ GeV}/c^2$  is investigated using a first-order model assuming weakly interacting  $b\bar{u}$  plus  $b\bar{d}$  meson clusters. The  $T_{bb}$  model yields a  $J^\pi = 1^+$  value in agreement with other analysis, and also suggests  $0^+$  and  $2^+$  possible assignments. Depending on the

assumed spin and parity of each meson cluster, the model yields mass values in the range of  $10.150 - 10.238 \text{ GeV}/c^2$ . These values are within about 4% of the proposed lattice QCD value.

**Keywords:**  $T_{bb}$  Tetraquark, First-Order Mass Formula, Quark Model, Cluster Model

#### 1. Introduction

As noted by Vujmilovic *et al.* [1], the postulated doubly-bottom tetraquark,  $T_{bb}$  with a  $bb\bar{u}\bar{d}$  has been scrutinized as one the most promising candidates for an exotic QCD stable state. Based on lattice QCD, Ref. 1 suggests that this state has a  $J^\pi = 1^+$  assignment with a mass of  $10.5765 \text{ GeV}/c^2$ . Advanced lattice studies determined that the  $T_{bb}$  mass is significantly below the lowest compatible decay threshold,  $BB^*$ . These studies are significant given the discovery of the  $T_{cc}$  with a  $cc\bar{u}\bar{d}$  configuration. The  $T_{bb}$  is investigated in terms of a first-order tetraquark model. This model is based on the semiempirical mass formula of Zel'dovich and Sakharov [2,3]. The first-order tetraquark model provides the  $T_{bb}$  mass as well its  $J^\pi$  value. Since the first-order model is limited in scope, it only permits a primitive angular momentum coupling structure. Other tetraquark systems [4-24] were reasonably described by the first-order model.

#### 2. Model and Formulation

The proposed first-order model is based on the semiempirical mass formula of Zel'dovich and Sakharov [2,3]. This model assumes that two weakly bound meson clusters form the tetraquark, with zero angular momentum between the clusters. The mesons ( $m$ ) mass ( $M$ ) is defined to have the form [2,3]:

$$\mathbf{M}_m = \delta_m + m_1 + m_2 + b_m [m_0^2 / (m_1 m_2)] \sigma_1 \cdot \sigma_2 \quad (1)$$

In Eq. 1,  $\delta_m$  is defined to have the value  $40 \text{ MeV}/c^2$  [3], and  $m_i$  is the mass of the quark comprising the meson cluster ( $i = 1$  and  $2$ ). The average mass of a first generation quark ( $u$  and  $d$ ) is  $m_0$  [25,26]. The scalar product of the quark spin vectors ( $\sigma_1 \cdot \sigma_2$ ) is  $-3/4$  and  $+1/4$  for pseudoscalar and vector mesons, respectively [3].

Effective quark masses provided by Griffiths [25] are used to determine the meson cluster mass. The  $d$ ,  $u$ ,  $s$ ,  $c$ ,  $b$ , and  $t$  quarks have effective mass values of 340, 336, 486, 1550, 4730, and 177000  $\text{MeV}/c^2$ , respectively [25]. Following the convention of the Standard Model, these quarks are grouped into three generations:  $[d(-1/3), u(+2/3)]$ ,  $[s(-1/3), c(+2/3)]$ , and  $[b(-1/3), t(+2/3)]$  [25,26]. The specific quark charges, in units of the unit charge  $e$ , are given within parentheses.

#### 3. First-Order Mass Formula for the $b\bar{u}$ plus $b\bar{d}$ Meson Cluster Description of the $T_{bb}$ Tetraquark

The first-order mass formula only provides a limited angular momentum coupling structure, and the spin of a tetraquark is derived from the angular momentum coupling of the two meson clusters:

$$\mathbf{J}^\pi = J^\pi(1) \times L \times J^\pi(2) \quad (2)$$

The first-order model summarized in Eq. 2 only provides a primitive angular momentum coupling structure for the  $J^\pi$  assignment, and the angular momentum between the clusters is zero ( $L = 0$ ). These are limiting conditions of the model. Detailed meson cluster structural information, and strong coupling between the clusters are not included in the model formulation.

These aforementioned simplifications minimize model complexity, and permit the tetraquark mass formula to have the form [5-24]:

$$M = M_m(1) + M_m(2) + \Phi \tag{3}$$

In Eq. 3, the two meson clusters are denoted by the numbers 1 and 2, and the individual meson cluster mass is given by Eq. 1. The final term in Eq. 3 ( $\Phi$ ) is the interaction between the meson clusters that is assumed to be negligible relative to the magnitude of the meson masses. Accordingly, Eq. 3 suggests a quasimolecular four quark system that is characteristic of a weakly bound meson-meson system.

The  $T_{bb}$  configuration is evaluated assuming weakly interacting  $b\bar{u}$  plus  $b\bar{d}$  meson clusters. The  $T_{bb}$  is modeled as the four possible configurations: a  $0^-$  and  $1^-$  ( $b\bar{d}$ ) meson cluster coupled to a  $0^-$  and  $1^-$  ( $b\bar{u}$ ) meson cluster. These assignments and their various couplings are provided in Table 1. The  $T_{bb}$  model yields a  $J^\pi = 1^+$  value in agreement with other analysis [1], and also suggests  $0^+$  and  $2^+$  possible assignments.

The predicted first-order mass is based on Eq. 3.

$$M(T_{bb}) = M(b\bar{d}) + M(b\bar{u}) + \Phi \tag{4}$$

Using Eq. 4 and the first-order mass formula of Eq. 1, mass values for the various meson cluster angular momentum configurations are provided in Table 1. Depending on the assumed spin and parity of each meson cluster, the model yields mass values in the range of 10.150 – 10.238 GeV/c<sup>2</sup>. These values are within about 4% of the proposed lattice QCD value.

**Table 1:** Model Results for the Proposed  $T_{bb}$  Tetraquark at 10.5765 GeV/c<sup>2</sup>

Meson Cluster - 1			Meson Cluster - 2			$T_{bb}$	
Configuration n	$J^\pi$	Mass (GeV/c <sup>2</sup> )	Configuration n	$J^\pi$	Mass (GeV/c <sup>2</sup> )	Mass (GeV/c <sup>2</sup> )	$J^\pi$
$b\bar{u}$	$0^-$	5.0728	$b\bar{d}$	$0^-$	5.0772	10.150	$0^+$
$b\bar{u}$	$0^-$	5.0728	$b\bar{d}$	$1^-$	5.1209	10.194	$1^+$
$b\bar{u}$	$1^-$	5.1171	$b\bar{d}$	$0^-$	5.0772	10.194	$1^+$
$b\bar{u}$	$1^-$	5.1171	$b\bar{d}$	$1^-$	5.1209	10.238	$0^+, 1^+, 2^+$

#### 4. First-Order Tetraquark Model Uncertainties and Limitations

There are a number of uncertainties and limitations that affect the model results. The limited angular momentum coupling structure restricts the available  $J^\pi$  values that can be evaluated. In addition, assuming zero angular momentum between the clusters also limits the evaluation of possible states.

The values for the effective quark masses [25] are not definitive. Although the weak coupling assumption appears to be reasonable, the exact magnitude for the interaction strength between the clusters is unknown [4-24]. The coupling strength will likely depend on the physical properties of the

interacting systems. In spite of these uncertainties, the model continues to provide reasonably credible results [4-24] for candidate tetraquark systems.

#### 5. Conclusions

The  $T_{bb}$  mass and  $J^\pi$  value are investigated using a first-order tetraquark model assuming weakly interacting  $b\bar{u}$  plus  $b\bar{d}$  meson clusters. The  $T_{bb}$  model yields a  $J^\pi = 1^+$  value in agreement with other analysis, and also suggests  $0^+$  and  $2^+$  possible assignments. Depending on the assumed spin and parity of each meson cluster, the model yields mass values in the range of 10.150 – 10.238 GeV/c<sup>2</sup>. These values are within about 4% of the proposed lattice QCD value.

#### 6. References

- Vujmilovic I, Collins S, Leskovec L, Prelovsek S. Electromagnetic form Factors and Structure of the  $T_{bb}$  Tetraquark from Lattice QCD, Phys. Rev. Lett. 2026; 136:161901.
- Ya B Zel'dovich, Sakharov AD. Kvarkovaia struktura i massy sil'novzaimodeistvuyushchikh chastits, Yad. Fiz. 1966; 4:395.
- Sakharov AD. Mass formula for mesons and baryons, Sov. Phys. JETP. 1980; 51:1059.
- Bevelacqua JJ. First-Order Tetraquark Mass Formula, Physics Essays. 2016; 29:198.
- Bevelacqua JJ. Description of the X(5568) and Proposed 750 GeV/c<sup>2</sup> State in Terms of a First-Order Tetraquark Mass Formula, Physics Essays. 2016; 29:367.
- Bevelacqua JJ. Fusion of Doubly Heavy Mesons into a Tetraquark, Physics Essays. 2018; 31:167.
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed X(3872), Physics Essays. 2019; 32:469.
- Bevelacqua JJ. Description of the X(6900) as a Four Charmed Quark State in Terms of a First-Order Tetraquark Mass Formula, Qeios **KLXLKJ**. 2020; 1. Doi: <https://doi.org/10.32388/KLXLKJ>
- Bevelacqua JJ. Description of the X(2900) as an Open Flavor Tetraquark in Terms of a First-Order Mass Formula, Qeios, **OVLMEB**. 2020; 1. Doi: <https://doi.org/10.32388/OVLMEB>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed  $Z_{cs}(3985)^-$ , Qeios **GLTEU2**. 2021; 1. Doi: <https://doi.org/10.32388/GLTEU2>
- Bevelacqua JJ. Possible Tetraquark Explanation for the X(6200), Qeios **J6AFYW**. 2021; 1. Doi: <https://doi.org/10.32388/J6AFYW>
- Bevelacqua JJ. Possible Tetraquark Explanation for the  $T_{cc}^+$ , Qeios **OMDGAQ**. 2021; 1. Doi: <https://doi.org/10.32388/OMDGAQ>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed  $Z_{cs}(4000)^+$  and  $Z_{cs}(4220)^+$ , Qeios **PPLMWV**. 2021; 1. Doi: <https://doi.org/10.32388/PPLMWV>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed X(3960), Qeios, **O1L0YM**. 2022; 1. Doi: <https://doi.org/10.32388/O1L0YM>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed  $T(2900)^{++}$  and  $T(2900)^0$  Structures, Qeios **V6WLTS**. 2022; 1. Doi: <https://doi.org/10.32388/V6WLTS>
- Bevelacqua JJ. Possible K K bar Tetraquark Explanation for the  $f_0(1370)$ , Qeios **HBDQXV**. 2023; 1. Doi: <https://doi.org/10.32388/HBDQXV>

17. Bevelacqua JJ. Possible f Quark Model of Tetraquarks and Pentaquarks, Qeios, **8T3IVE**. 2023; 1. Doi: <https://doi.org/10.32388/8T3IVE>
18. Bevelacqua JJ. Possible Tetraquark Explanation for the  $\Upsilon(10753)$ , Qeios **NZRGH3**. 2023; 1. Doi: <https://doi.org/10.32388/NZRGH3>
19. Bevelacqua JJ. Possible Tetraquark Explanation for the  $\psi(4230)$ ,  $\psi(4360)$ , and  $\psi(4415)$ , Qeios **D5HKO0**. 2024; 1. Doi: <https://doi.org/10.32388/D5HKO0>
20. Bevelacqua JJ. Possible Tetraquark Explanation for the  $J/\psi K_s^0$  Structure Observed in Proton-Proton Collision Data at Center-of-Mass Energies of 7, 8, and 13 TeV, Qeios **YFT8L5**. 2024; 1. Doi: <https://doi.org/10.32388/YFT8L5>
21. Bevelacqua JJ. Possible Explanations for the Proposed  $J/\Psi + J/\Psi$  and  $J/\Psi + \Psi(2S)$  Tetraquark States, Qeios **W2A4LD**. 2024; 1. Doi: <https://doi.org/10.32388/W2A4LD>
22. Bevelacqua JJ. Possible Tetraquark Explanations for the B D-bar and B-star D-bar States Proposed by Lattice QCD Calculations, Qeios **8MWBEY**. 2024; 1. Doi: <https://doi.org/10.32388/8MWBEY>
23. Bevelacqua JJ. Possible Tetraquark Explanations for the R(3780), Qeios **WRVP7E**. 2025; 1. Doi: <https://doi.org/10.32388/WRVP7E>
24. Bevelacqua JJ. Possible  $\omega$   $\Upsilon(1S)$  Tetraquark Explanation for the  $\chi_{b0}(2P)$ . Int. J. Adv. Multidisc. Res. Stud. 2025; 5(6):35.
25. Griffiths D. Introduction to Elementary Particles, 2<sup>nd</sup> ed., Wiley-VCH, Weinheim, 2008.
26. Particle Data Group, Review of Particle Physics, Phys. Rev. D. 2024; 110:030001.