



Received: 18-05-2026
Accepted: 28-06-2026

ISSN: 2583-049X

Proposed T_{bc} Tetraquark Near the $B^*\bar{D}$ Threshold

JJ Bevelacqua

Bevelacqua Resources, 7531 Flint Crossing Circle SE, Owens Cross Roads, AL 35763, USA

Corresponding Author: JJ Bevelacqua

Abstract

The proposed T_{bc} tetraquark near the $B^*\bar{D}$ threshold is investigated using a first-order model assuming weakly interacting $b\bar{u}$ plus $c\bar{d}$ meson clusters. The T_{bc} model yields a $J^\pi = 0^+$ and 1^+ values in agreement with other analysis, and also suggests a 2^+ possible assignment. Depending on the

assumed spin and parity of each meson cluster, the model yields mass values in the range of 6.9028 – 7.0804 GeV/c². These values are in reasonable agreement of 7.0944 – 7.1514 GeV/c² with proposed lattice QCD values.

Keywords: T_{bc} Tetraquark, First-Order Mass Formula, Quark Model, Cluster Model

1. Introduction

Alexandrou *et al.* [1] observed sharp peaks in the $B - \bar{D}$ and $B^* - \bar{D}$ scattering rates close to the thresholds. These resonances are associated with shallow bound states, a few MeV or less below these thresholds. In addition, Ref. 1 found hints for resonances with masses of order 100 MeV above these thresholds. In Ref. 2 these states were assigned possible $J^\pi = 0^+$ and 1^+ values. In addition, Ref. 3 reported a $J^\pi = 1^+ b\bar{c}ud$ tetraquark bound states using lattice QCD 43 MeV below the $B^* - \bar{D}$ threshold.

These results suggest possible a T_{bc} tetraquark with a mass between 7.0944 and 7.1514 GeV/c² with possible $J^\pi = 0^+$ and 1^+ values. Definitive experimental results have not yet been published.

The T_{bc} is investigated in terms of a first-order tetraquark model. This model is based on the semiempirical mass formula of Zel'dovich and Sakharov [4, 5]. The first-order tetraquark model provides the T_{bc} mass as well its J^π value. Since the first-order model is limited in scope, it only permits a primitive angular momentum coupling structure. Other tetraquark systems [6-27] were reasonably described by the first-order model.

2. Model and Formulation

The proposed first-order model is based on the semiempirical mass formula of Zel'dovich and Sakharov [4, 5]. This model assumes that two weakly bound meson clusters form the tetraquark, with zero angular momentum between the clusters.

The mesons (m) mass (M) is defined to have the form [4, 5]:

$$\mathbf{M}_m = \delta_m + \mathbf{m}_1 + \mathbf{m}_2 + \mathbf{b}_m [\mathbf{m}_0^2 / (\mathbf{m}_1 \mathbf{m}_2)] \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 \quad (1)$$

In Eq. 1, δ_m is defined to have the value 40 MeV/c² [5], and m_i is the mass of the quark comprising the meson cluster ($i = 1$ and 2). The average mass of a first generation quark (u and d) is m_0 [28, 29]. The scalar product of the quark spin vectors ($\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2$) is $-3/4$ and $+1/4$ for pseudoscalar and vector mesons, respectively [5].

Effective quark masses provided by Griffiths [28] are used to determine the meson cluster mass. The d , u , s , c , b , and t quarks have effective mass values of 340, 336, 486, 1550, 4730, and 177000 MeV/c², respectively [28]. Following the convention of the Standard Model, these quarks are grouped into three generations: [$d(-1/3)$, $u(+2/3)$], [$s(-1/3)$, $c(+2/3)$], and [$b(-1/3)$, $t(+2/3)$] [28, 29]. The specific quark charges, in units of the unit charge e , are given within parentheses.

3. First-Order Mass Formula for the $b\bar{u}$ plus $c\bar{d}$ Meson Cluster Description of the T_{bc} Tetraquark

The first-order mass formula only provides a limited angular momentum coupling structure, and the spin of a tetraquark is derived from the angular momentum coupling of the two meson clusters:

$$\mathbf{J}^\pi = \mathbf{J}^\pi(1) \times \mathbf{L} \times \mathbf{J}^\pi(2) \tag{2}$$

The first-order model summarized in Eq. 2 only provides a primitive angular momentum coupling structure for the J^π assignment, and the angular momentum between the clusters is zero ($L = 0$). These are limiting conditions of the model. Detailed meson cluster structural information, and strong coupling between the clusters are not included in the model formulation.

These aforementioned simplifications minimize model complexity, and permit the tetraquark mass formula to have the form [6-27]:

$$\mathbf{M} = \mathbf{M}_m(1) + \mathbf{M}_m(2) + \Phi \tag{3}$$

In Eq. 3, the two meson clusters are denoted by the numbers 1 and 2, and the individual meson cluster mass is given by Eq. 1. The final term in Eq. 3 (Φ) is the interaction between the meson clusters that is assumed to be negligible relative to the magnitude of the meson masses. Accordingly, Eq. 3 suggests a quasimolecular four quark system that is

characteristic of a weakly bound meson-meson system.

The T_{bc} configuration is evaluated assuming weakly interacting $b\bar{u}$ plus $c\bar{d}$ meson clusters. The T_{bc} is modeled as the four possible configurations: a 0^- and 1^- ($c\bar{d}$) meson cluster coupled to a 0^- and 1^- ($b\bar{u}$) meson cluster. These assignments and their various couplings are provided in Table 1. The T_{bc} model yields a $J^\pi = 1^+$ value in agreement with other analysis [1], and also suggests 0^+ and 2^+ possible assignments.

The predicted first-order mass is based on Eq. 3:

$$\mathbf{M}(T_{bc}) = \mathbf{M}(c\bar{d}) + \mathbf{M}(b\bar{u}) + \Phi \tag{4}$$

Using Eq. 4 and the first-order mass formula of Eq. 1, mass values for the various meson cluster angular momentum configurations are provided in Table 1. Depending on the assumed spin and parity of each meson cluster, the model yields mass values in the range of 6.9028– 7.0804 GeV/c². These values are in reasonable agreement with proposed 7.0944 – 7.1514 GeV/c² lattice QCD values [1-3].

Table 1: Model Results for the Proposed T_{bc} Tetraquark near the $B^*\bar{D}$ threshold

Meson Cluster - 1			Meson Cluster - 2			T_{bc}	
Configuration	J^π	Mass (GeV/c ²)	Configuration	J^π	Mass (GeV/c ²)	Mass (GeV/c ²)	J^π
$b\bar{u}$	0^-	5.0728	$c\bar{d}$	0^-	1.8300	6.9028	0^+
$b\bar{u}$	0^-	5.0728	$c\bar{d}$	1^-	1.9633	7.0361	1^+
$b\bar{u}$	1^-	5.1171	$c\bar{d}$	0^-	1.8300	6.9471	1^+
$b\bar{u}$	1^-	5.1171	$c\bar{d}$	1^-	1.9633	7.0804	$0^+, 1^+, 2^+$

4. First-Order Tetraquark Model Uncertainties and Limitations

There are a number of uncertainties and limitations that affect the model results. The limited angular momentum coupling structure restricts the available J^π values that can be evaluated. In addition, assuming zero angular momentum between the clusters also limits the evaluation of possible states.

The values for the effective quark masses [28] are not definitive. Although the weak coupling assumption appears to be reasonable, the exact magnitude for the interaction strength between the clusters is unknown [6-27]. The coupling strength will likely depend on the physical properties of the interacting systems. In spite of these uncertainties, the model continues to provide reasonably credible results [6-27] for candidate tetraquark systems.

5. Justification for the Weak Coupling Assumption

The quark charges are related to the number of colors (N_c) incorporated into the fundamental quantum chromodynamics (QCD) formulation [30, 31]. For example, the first generation quark charges within $SU(N_c)$ are:

$$\mathbf{Q}_d = \frac{1}{2} \left(\frac{1}{N_c} - 1 \right) e \tag{5}$$

$$\mathbf{Q}_u = \frac{1}{2} \left(\frac{1}{N_c} + 1 \right) e \tag{6}$$

For conventional QCD using 3 colors, the expected d and u electric charges are obtained. The importance of QCD expansions involving $1/N_c$ illustrates the weak coupling

assumption [32, 33].

A key assumption of the first-order mass formula of Eqs. 1, 3, and 4 is weak coupling between the two clusters. In particular, the model utilized in this paper assumes the tetraquark is partitioned into two clusters with the interaction between the clusters providing a minimal contribution to the tetraquark mass. Within the scope of this mass formula, the meson-meson cluster interaction is assumed in Eq. 4 to be weak and sufficiently small to be ignored.

This assumption is justified because QCD can be investigated as an expansion in $1/N_c$ [32, 33]. The large N_c limit reduces to a field theory of weakly interacting meson-like objects. The physical situation with $N_c = 3$ retains many of the characteristics of the $N_c \rightarrow \infty$ limit, and further justifies the weak coupling approximation.

The $1/N_c$ expansion [32, 33] is well accepted in elementary particle physics and leads to the Okubo-Zweig-Iizuka (OZI) rule [34-36] and the Skyrme model [37, 38]. In fact, in the $1/N_c$ expansion, QCD is reduced to a weakly interacting meson theory, and the meson-meson interaction is regarded to be small [32, 33]. This situation is also a characteristic of Eqs. 3 and 4.

6. Conclusions

The proposed T_{bc} tetraquark near the $B^*\bar{D}$ threshold is investigated using a first-order model assuming weakly interacting $b\bar{u}$ plus $c\bar{d}$ meson clusters. This weak coupling approximation is based on the $1/N_c$ expansion that is well accepted in elementary particle physics and leads to the Okubo-Zweig-Iizuka (OZI) rule and the Skyrme model. In addition, in the $1/N_c$ expansion, QCD is reduced to a weakly interacting meson theory, and the meson-meson interaction

is regarded to be small. This situation is also a characteristic of Eqs. 3 and 4.

The T_{bb} model yields a $J^{\pi} = 0^{+}$ and 1^{+} values in agreement with other analysis, and also suggests a 2^{+} possible assignment. Depending on the assumed spin and parity of each meson cluster, the model yields mass values in the range of $6.9028 - 7.0804 \text{ GeV}/c^2$. These values are in reasonable agreement with proposed $7.0944 - 7.1514 \text{ GeV}/c^2$ lattice QCD values.

7. References

- Alexandrou C, *et al.* Shallow Bound States and Hints for Broad Resonances with Quark Content $b\bar{c}u\bar{d}$ in $B - \bar{D}$ and $B^{*} - \bar{D}$ Scattering from Lattice QCD, *Phys. Rev. Lett.* 2024; 132:151902.
- Meinel S, Pflaumer M, Wagner M. Search for $\bar{b}b\bar{u}s$ and $b\bar{c}u\bar{d}$ tetraquark bound states using lattice QCD, *Phys. Rev. D.* 2022; 106:034507.
- Padmanath M, Radhakrishnan A, Mathur N. Bound isoscalar axial-vector $b\bar{c}u\bar{d}$ tetraquark T_{bc} in QCD, arXiv:2307.14128.v2 [hep-lat], May 1, 2024.
- Ya B Zel'dovich, Sakharov AD. Kvarkovaia struktura i massy sil'novzaimodeistvuyushchikh chastits, *Yad. Fiz.* 1966; 4:395.
- Sakharov AD. Mass formula for mesons and baryons, *Sov. Phys. JETP.* 1980; 51:1059.
- Bevelacqua JJ. First-Order Tetraquark Mass Formula, *Physics Essays.* 2016; 29:198.
- Bevelacqua JJ. Description of the $X(5568)$ and Proposed $750 \text{ GeV}/c^2$ State in Terms of a First-Order Tetraquark Mass Formula, *Physics Essays.* 2016; 29:367.
- Bevelacqua JJ. Fusion of Doubly Heavy Mesons into a Tetraquark, *Physics Essays.* 2018; 31:167.
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed $X(3872)$, *Physics Essays.* 2019; 32:469.
- Bevelacqua JJ. Description of the $X(6900)$ as a Four Charmed Quark State in Terms of a First-Order Tetraquark Mass Formula, *Qeios* **KLXLKJ**. 2020; 1. Doi: <https://doi.org/10.32388/KLXLKJ>
- Bevelacqua JJ. Description of the $X(2900)$ as an Open Flavor Tetraquark in Terms of a First-Order Mass Formula, *Qeios*, **OVLMEB**. 2020; 1. Doi: <https://doi.org/10.32388/OVLMEB>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed $Z_{cs}(3985)^{-}$, *Qeios* **GLTEU2**. 2021; 1. Doi: <https://doi.org/10.32388/GLTEU2>
- Bevelacqua JJ. Possible Tetraquark Explanation for the $X(6200)$, *Qeios* **J6AFYW**. 2021; 1. Doi: <https://doi.org/10.32388/J6AFYW>
- Bevelacqua JJ. Possible Tetraquark Explanation for the T_{cc}^{+} , *Qeios* **OMDGAQ**. 2021; 1. Doi: <https://doi.org/10.32388/OMDGAQ>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed $Z_{cs}(4000)^{+}$ and $Z_{cs}(4220)^{+}$, *Qeios* **PPLMWV**. 2021; 1. Doi: <https://doi.org/10.32388/PPLMWV>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed $X(3960)$, *Qeios*, **O1L0YM**. 2022; 1. Doi: <https://doi.org/10.32388/O1L0YM>
- Bevelacqua JJ. Possible Tetraquark Explanation for the Proposed $T(2900)^{++}$ and $T(2900)^0$ Structures, *Qeios* **V6WLTS**. 2022; 1. Doi: <https://doi.org/10.32388/V6WLTS>
- Bevelacqua JJ. Possible $K K \bar{K}$ Tetraquark Explanation for the $f_0(1370)$, *Qeios* **HBDQXV**. 2023; 1. Doi: <https://doi.org/10.32388/HBDQXV>
- Bevelacqua JJ. Possible f Quark Model of Tetraquarks and Pentaquarks, *Qeios*, **8T3IVE**. 2023; 1. Doi: <https://doi.org/10.32388/8T3IVE>
- Bevelacqua JJ. Possible Tetraquark Explanation for the $Y(10753)$, *Qeios* **NZRGH3**. 2023; 1. Doi: <https://doi.org/10.32388/NZRGH3>
- Bevelacqua JJ. Possible Tetraquark Explanation for the $\psi(4230)$, $\psi(4360)$, and $\psi(4415)$, *Qeios* **D5HKO0**. 2024; 1. Doi: <https://doi.org/10.32388/D5HKO0>
- Bevelacqua JJ. Possible Tetraquark Explanation for the $J/\psi K^0_s$ Structure Observed in Proton-Proton Collision Data at Center-of-Mass Energies of 7, 8, and 13 TeV, *Qeios* **YFT8L5**. 2024; 1. Doi: <https://doi.org/10.32388/YFT8L5>
- Bevelacqua JJ. Possible Explanations for the Proposed $J/\Psi + J/\Psi$ and $J/\Psi + \Psi(2S)$ Tetraquark States, *Qeios* **W2A4LD**. 2024; 1. Doi: <https://doi.org/10.32388/W2A4LD>
- Bevelacqua JJ. Possible Tetraquark Explanations for the $B D\text{-bar}$ and $B\text{-star } D\text{-bar}$ States Proposed by Lattice QCD Calculations, *Qeios* **8MWBEY**. 2024; 1. Doi: <https://doi.org/10.32388/8MWBEY>
- Bevelacqua JJ. Possible Tetraquark Explanations for the $R(3780)$, *Qeios* **WRVP7E**. 2025; 1. Doi: <https://doi.org/10.32388/WRVP7E>
- Bevelacqua JJ. Possible $\omega \text{ } Y(1S)$ Tetraquark Explanation for the $\chi_{b0}(2P)$. *Int. J. Adv. Multidisc. Res. Stud.* 2025; 5(6):35.
- Bevelacqua JJ. Proposed T_{bb} Tetraquark at $10.5765 \text{ GeV}/c^2$. *Int. J. Adv. Multidisc. Res. Stud.* 2026; 6(3):635.
- Griffiths D. Introduction to Elementary Particles, 2nd ed., Wiley-VCH, Weinheim, 2008.
- Particle Data Group, Review of Particle Physics, *Phys. Rev. D.* 2024; 110:030001.
- Abbas A. Anomalies and charge quantization in the standard model with arbitrary number of colors, *Phys. Lett. B.* 1990; 238:344.
- Abbas A. Spontaneous symmetry breaking, quantization of the electric charge and the anomalies, *J. Phys. G.* 1990; 16:L163.
- 't Hooft G. A Planar Diagram Theory for Strong Interactions, *Nucl. Phys. B.* 1974; 72:461.
- Witten E. Baryons in the $1/N$ Expansion, *Nucl. Phys. B.* 1979; 160:57.
- Okubo S. ϕ -meson and unitary symmetry model, *Phys. Lett.* 1963; 5:1975.
- Zweig G. An $SU(3)$ model for strong interaction symmetry and its breaking-II, CERN Report No. 8419/TH412, 1964.
- Iizuka J. A Systematics and Phenomenology of Meson Family, *Prog. Theor. Phys. Suppl.* 1966; 21:37-38.
- Skyrme THR. A non-linear field theory, *Proc. R. Soc. A.* 1961; 260:127.
- Skyrme THR. A unified field theory of mesons and baryons, *Nucl. Phys.* 1962; 31:556.