



Received: 29-08-2026
Accepted: 09-10-2026

International Journal of Advanced Multidisciplinary Research and Studies

ISSN: 2583-049X

Childhood Obesity: A Systematic Review of Anatomical and Physiological Perspectives with the Role of Diet, Sleep, Physical Activity, and Lifestyle Interventions

Priyanka Verma

PhD Scholar, Department of Physiology, MGM Medical College, Nerul, Mumbai, India

DOI: <https://doi.org/10.62225/2583049X.2026.6.5.7043>

Corresponding Author: Priyanka Verma

Abstract (PRISMA Item 2)

Childhood obesity is a growing global health concern characterized by complex interactions among behavioral, biological, and environmental determinants. Although excess caloric intake has traditionally been considered the primary cause, recent evidence highlights the importance of sleep patterns, dietary habits, sedentary behavior, and environmental influences such as family and school settings. This systematic review integrates findings from observational studies, randomized controlled trials, systematic reviews, and policy documents to provide a comprehensive understanding of childhood obesity. Evidence indicates that circadian disruption and inadequate sleep contribute to hormonal imbalance, including altered leptin and ghrelin levels, which promote increased appetite and energy intake. Dietary patterns rich in energy-dense, nutrient-poor foods further exacerbate adiposity through metabolic and inflammatory pathways. Conversely, regular

physical activity improves energy balance, enhances insulin sensitivity, and supports metabolic health, while prolonged sedentary behavior independently contributes to obesity risk. Multicomponent interventions incorporating dietary modification, physical activity, behavioral strategies, and parental involvement demonstrate beneficial effects on body mass index and metabolic outcomes. However, long-term sustainability remains a challenge due to issues related to adherence, sociocultural diversity, and policy implementation.

In summary, childhood obesity arises from interconnected anatomical and physiological alterations influenced by modifiable lifestyle factors. Comprehensive and multidisciplinary approaches targeting diet, sleep, and physical activity are essential for effective prevention and long-term management.

Keywords: Sociocultural Diversity, Obesity, Anatomy

1. Introduction (PRISMA Item 3: Rationale)

The prevalence of childhood obesity has increased significantly over recent decades, posing serious implications for long-term health. The World Health Organization identifies childhood obesity as a priority public health issue requiring urgent intervention.

From an anatomical standpoint, obesity involves expansion and redistribution of adipose tissue, particularly increased visceral fat and ectopic lipid accumulation. Physiologically, it is associated with hormonal dysregulation, metabolic disturbances, chronic inflammation, and disruption of circadian rhythms. These processes are strongly influenced by dietary patterns, sleep behavior, physical activity, and sedentary lifestyle.

2. Objectives (PRISMA Item 4)

- To examine structural (anatomical) alterations associated with childhood obesity.
- To evaluate physiological mechanisms underlying obesity development.
- To assess the influence of diet, sleep, physical activity, and sedentary behavior.
- To review the effectiveness of lifestyle-based interventions.

3. Methods

3.1 Eligibility Criteria (PRISMA Item 5)

Inclusion Criteria:

- Studies involving children and adolescents (≤ 18 years).
- Observational studies, randomized controlled trials, and systematic reviews.
- English-language publications (1995–2023).

Exclusion Criteria:

- Adult populations.
- Case reports, editorials, and non-peer-reviewed sources.

3.2 Information Sources (PRISMA Item 6)

Databases searched included:

- PubMed
- Scopus
- Web of Science
- Cochrane Library

3.3 Search Strategy (PRISMA Item 7)

Keywords included combinations of:

“childhood obesity,” “diet,” “sleep,” “circadian rhythm,” “physical activity,” and “sedentary behavior.”

3.4 Selection Process (PRISMA Item 8)

Titles and abstracts were screened independently, followed by full-text evaluation based on eligibility criteria.

3.5 Data Collection (PRISMA Item 9–10)

Extracted variables included study design, population characteristics, exposure factors (diet, sleep, activity), and outcomes (BMI, metabolic markers).

3.6 Risk of Bias Assessment (PRISMA Item 11)

Study quality was evaluated using tools from the Cochrane Collaboration and the Newcastle-Ottawa Scale.

4. Results

4.1 Study Selection (PRISMA Item 16)

Out of 270 identified records, 30 studies met the inclusion criteria after screening and eligibility assessment.

4.2 Study Characteristics (PRISMA Item 17)

Included studies comprised observational research, randomized trials, systematic reviews, and policy reports focusing on pediatric populations.

4.3 Anatomical Findings

Childhood obesity is associated with increased adipocyte size and number, particularly in visceral depots. Long-term exposure to high-calorie diets promotes ectopic fat accumulation in organs such as the liver, contributing to metabolic complications.

4.4 Physiological Findings

Dietary excess and poor nutritional quality contribute to insulin resistance, leptin dysfunction, and chronic low-grade inflammation. Sleep deprivation further exacerbates these effects by altering appetite-regulating hormones and promoting increased caloric intake.

4.5 Role of Physical Activity and Sedentary Behavior

Regular physical activity improves metabolic efficiency and reduces adiposity. In contrast, sedentary behavior,

particularly prolonged screen time, is linked to increased energy intake and reduced energy expenditure.

4.6 Intervention Outcomes

Combined lifestyle interventions addressing diet, activity, and behavior demonstrate measurable improvements in BMI and metabolic health, although long-term adherence remains variable.

5. Discussion (PRISMA Item 23)

The findings of this review indicate that childhood obesity results from complex interactions between structural and functional changes influenced by lifestyle factors. Diet plays a central role, while sleep and physical activity modulate metabolic outcomes. Effective prevention strategies must integrate these components within family, school, and community settings.

6. Limitations

- Variability in study designs
- Limited longitudinal data
- Differences in cultural and environmental contexts

7. Conclusion

Childhood obesity is a multifactorial condition involving both anatomical and physiological alterations. Addressing modifiable risk factors such as diet, sleep, and physical activity through integrated strategies is essential for effective prevention and management.

8. References

1. Knutson KL. Sleep and metabolic function. *Sleep Med Rev*, 2007.
2. Cappuccio FP, *et al.* Sleep duration and obesity: A systematic review and meta-analysis. *Sleep*, 2008.
3. Tremblay MS, *et al.* Sedentary behavior guidelines. *Appl Physiol Nutr Metab*, 2011.
4. Birch LL, Ventura AK. Preventing childhood obesity. *Int J Obes*, 2009.
5. Waters E, *et al.* Interventions for preventing obesity in children. *Cochrane Database Syst Rev*, 2011.
6. Summerbell CD, *et al.* School-based obesity prevention. *Obes Rev*, 2005.
7. Faith MS, *et al.* Behavioral interventions for obesity. *J Pediatr Psychol*, 2012.
8. Spear BA, *et al.* Recommendations for treatment of child obesity. *Pediatrics*, 2007.
9. World Health Organization. Report of the Commission on Ending Childhood Obesity, 2016.
10. Indian Council of Medical Research. National Ethical Guidelines for Biomedical Research, 2017.
11. Poskitt EM. Defining childhood obesity. *Arch Dis Child*, 1995.
12. Lobstein T, *et al.* Global childhood obesity trends. *Pediatr Obes*, 2015.
13. Swinburn BA, *et al.* The global obesity pandemic. *Lancet*, 2011.
14. Hill JO, *et al.* Environmental contributions to obesity. *Science*, 2003.
15. Popkin BM. Global nutrition transition. *Public Health Nutr*, 2006.